

HANDBOOK OF THE ELEMENTS

IA											
H	IIA										
Li	Be										
Na	Mg	IIIB	IVB	VB	VIB	VII B	VIII			IB	II B
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd
Cs	Ba	La*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg
Fr	Ra	Act†	104	105	106	107		109			
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md

SAMUEL RUBEN

											O								
						III A	IV A	V A	VI A	VII A	He								
						B	C	N	O	F	Ne								
						Al	Si	P	S	Cl	Ar								
VII B		VIII			IB	IIB													
Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr								
Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe								
Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn								
107		109																	
<table border="1"> <tr> <td>Gd</td> <td>Tb</td> <td>Dy</td> <td>Ho</td> <td>Er</td> <td>Tm</td> <td>Yb</td> <td>Lu</td> </tr> </table>												Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu												
<table border="1"> <tr> <td>Cm</td> <td>Bk</td> <td>Cf</td> <td>Es</td> <td>Fm</td> <td>Md</td> <td>No</td> <td>Lr</td> </tr> </table>												Cm	Bk	Cf	Es	Fm	Md	No	Lr
Cm	Bk	Cf	Es	Fm	Md	No	Lr												

OPEN COURT
LA SALLE, ILLINOIS 61301
ISBN 0-87548-399-2

\$10.95

ISBN 0-87548-399-2



9 780875 483993

Handbook of the Elements

Handbook of the Elements

Samuel Ruben

**Open Court Publishing Company
La Salle, Illinois 61301**

**To order books from Open Court,
call toll-free 1-800-815-2280.**

Open Court Publishing Company is a division of Carus
Publishing Company.

Copyright © 1965, 1967, 1985 by Samuel Ruben.

Fourth printing 1992

Fifth printing 1994

Sixth printing 1995

Seventh printing 1996

Eighth printing 1998

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher, Open Court Publishing Company, a division of Carus Publishing Company, 315 Fifth Street, P.O. Box 300, Peru, Illinois 61354-0300.

Printed and bound in the United States of America.

Library of Congress Cataloging-in-Publication Data

Ruben, Samuel

Handbook of the elements.

1. Chemical elements. I. Title

QD466.R78 1985 546 85-18942

ISBN 0-87548-399-2

Preface

Handbook of the Elements is a practical reference source that provides essential information on the 108 known chemical elements for students and working scientists alike.

Knowledge about the elements is critical to our understanding of science and the world around us. This edition represents the most up-to-date compilation of information on the elements currently available.

Data on the chemical elements have been the fundamentals of scientific work for years, yet new research is continually revising previously published material about them. Even physical "constants" are subject to change in the light of additional research.


The information contained in this the third edition reflects state-of-the-art values on the most frequently required constants. The material in this current edition was compiled, corrected, and updated over a period of several years, utilizing hundreds of sources. Each value was checked in a minimum of 10 sources to ensure accuracy. A partial listing of the primary reference sources consulted is given at the end of the monographs.

I wish to acknowledge the significant assistance of Wayne Hruden for updating the reported values of the constants and the support given by the Duracell International Inc.

SAMUEL RUBEN
December 1984

Introduction

This handbook contains monographs for each of the 108 known chemical elements, arranged in alphabetical order for rapid reference.



Except where unavailable, values for the following twenty-five different elemental constants are given:

Periodic classification The group, family name, and/or series of the element; this categorization reflects the position of the element in the periodic table.

Atomic number An element of atomic number Z occupies the Z th position in the periodic classification. Its neutral atom has a nucleus with a charge of $+Ze$ surrounded by Z electrons, each of charge $-e$.

Atomic weight The relative atomic mass (A_r) based on $^{12}\text{C} \equiv 12$; the value for the most stable isotope is given for synthetic elements.

-
- Naturally occurring isotopes** Mass numbers of the isotopes are listed in decreasing order of natural terrestrial abundance.
- Density** The weight per unit volume of the element; measurements of this constant are generally made at 25°C, but the temperature utilized is shown in parentheses. Units are grams per cubic centimeter (g/cm^3).
- Melting point** Units are degrees Celsius ($^{\circ}\text{C}$); **Boiling point:** Units are degrees Celsius ($^{\circ}\text{C}$).
- Latent heat of fusion** The quantity of heat required to change 1 g of the solid element into the liquid state at a constant temperature. Units are Joules per gram (J/g).
- Specific heat** The thermal capacity of an element; the specific heat capacity is the quantity of heat required to raise the temperature of a mass through a measured number of Celsius degrees. Units are Joules per gram per degree Celsius ($\text{J}/\text{g}^{\circ}\text{C}$).
- Coefficient of lineal thermal expansion** The ratio of the change in length per degree Celsius to the original length at zero degrees Celsius. Units are centimeter per centimeter per degree Celsius ($\text{cm}/\text{cm}^{\circ}\text{C}$).
- Thermal conductivity** Thermal energy transmitted through a unit cube per unit time when there exists unit temperature difference between opposite parallel faces. Units are watts (or milliwatts) per centimeter per degree Celsius [w (or mw)/ $\text{cm}^{\circ}\text{C}$].
- Electrical resistivity** A proportionality factor (ρ) relating the resistance to current flow between parallel faces of a 1-cm cube of the element. This factor is also known as specific resistance. Because the resistance of semiconductor is substantially influenced by the presence of traces of impurities, the intrinsic resistivity is the parameter given for these ultrapure elements. Units are ohm-centimeters (ohm-cm).
- Ionization potential (1st)** The energy necessary to remove the least strongly bound electron from its orbit and place it at rest at an infinite distance. Units are electron volts (eV).
- Electron work function (ϕ)** The minimum photonic energy required to remove an electron from the boundary of an element; also known as photoelectric work function. Units are electron volts (eV).
- Oxidation potential** The difference in potential produced by a voltaic half-cell associated with the cited chemical reaction. By using the oxidation potential, the likelihood of various chemical reactions can be predicted. Oxidation of gaseous hydrogen (at 1 atmosphere pressure) to ionic hydrogen (in 1 molar acid solution at 25°C) defines the zero reference. Units are volts (V).
- Chemical valence** The number of hydrogen atoms (or their equivalent) with which an atom of an element can combine (if negative) or the number which it can displace in a reaction (if positive). The principal valence is set in italic type when more than one valence is possible.

Electrochemical equivalents The mass of an element displaced by the passage of unit quantity of electricity. The values provided are derived from:

$$\text{electrochemical equivalents} = \frac{kA}{n}$$

where k is a constant equal to 0.0373100, A is the gram-atomic weight, and n is the principal valence. Units are grams per ampere-hour (g/amp-hr).

Ionic radius The radius an ion exhibits in an ionic crystal in which the ions are packed together with their outermost electronic shells in contact with each other. Values are given for a coordination number of 6. Ionic radii for other coordination numbers can be obtained by multiplying by the following conversion factors:

<i>Coordination Number</i>	<i>Conversion Factor</i>
12	1.12
9	1.05
8	1.03
6	1.00
4	0.94

Units are Ångstroms ($1\text{Å} = 10^{-8}\text{ cm}$).

Valence electron potential ($-\epsilon V$) A calculated value based on the charge of the valence electrons and the ionic radius. It provides a quantitative indication of the reactivity of an element and is determined by the equation:

$$(-\epsilon V) = \frac{kn}{r}$$

where $(-\epsilon V)$ is the valence electron potential, n is the valence, and k is a proportionality factor converting Ångstroms to centimeters and expressing the force exerted by the valence electrons in electron volts and is equal to 14.399; r is the ionic radius in Ångstroms. The principal valence has been used for the determination.

Electronic configuration A sequential listing of the orbiting electrons, indicating the principal shells and the number of electrons in each subshell. For example, $4d^{10}$ would indicate the presence of 10 electrons in the "d" subshell of the fourth (N) principal shell. Principal shells are assigned letters corresponding to their quantum numbers as follows: 1 = K, 2 = L, 3 = M, 4 = N, 5 = O, 6 = P, and 7 = Q. A maximum exists for the number of electrons in each subshell: 2 in s, 6 in p, 10 in d, and 14 in f.

Valence electrons A sequential listing of the electrons involved in the ionization of the element. They are indicated in the same manner as in the electronic configuration.


Crystal form A brief description of the atomic arrangement in the elemental solid state. (See accompanying figure for common Crystal Forms).

Half life The time required for one-half of an initial quantity of a radioactive isotope to be converted into its decay product. This entry is included only when all known isotopes of an element are unstable. The half life presented is that of the most stable isotope. Units are seconds, minutes, hours, days, or years.


Cross section σ The effective size of a nucleus in capturing a thermal (slow) neutron. The larger the cross section the greater is the probability of neutron capture. Units are millibarns (mbarns) or barns (1 barn = 10^{-24} cm²).

Vapor pressure The pressure exerted when a solid or liquid is in equilibrium with its vapor. Since this parameter is a function of temperature, the vapor pressure at the melting point is given. Units are Pascals (Pa).


Crystal Forms




CUBIC, BODY CENTERED




CUBIC, FACE CENTERED




CUBIC, DIAMOND




RHOMBOHEDRAL
 $\alpha, \beta, \gamma, \neq 90^\circ$




TETRAGONAL
 $a \neq c$



MONOCLINIC
 $\alpha \neq 90^\circ$



ORTHORHOMBIC
 $a \neq b \neq c$



HEXAGONAL
 $a \neq c$

Periodic Classification of the Elements

IA										
1.0079 H 1	IIA									
6.941 Li 3	9.01218 Be 4									
22.98977 Na 11	24.305 Mg 12				IIIB	IVB	VB	VIB	VIIB	VIII
39.098 K 19	40.08 Ca 20	44.95592 Sc 21	47.90 Ti 22	50.9415 V 23	51.996 Cr 24	54.9380 Mn 25	55.847 Fe 26	58.9332 Co 27	58.70 Ni 28	
85.4678 Rb 37	87.62 Sr 38	88.9059 Y 39	91.22 Zr 40	92.9064 Nb 41	95.94 Mo 42	96.906 Tc 43	101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46	
132.9054 Cs 55	137.34 Ba 56	138.9055 La* 57	178.49 Hf 72	180.9479 Ta 73	183.85 W 74	186.2 Re 75	190.2 Os 76	192.22 Ir 77	195.09 Pt 78	
223.01976 Fr 87	226.02544 Ra 88	227.02777 Act 89	104	105	106	107		109		

*Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
--------------------	----------------------	--------------------	---------------------	-------------------	--------------------	--------------------	----------------------	--------------------	----------------------

†Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
-----------------------	----------------------	--------------------	----------------------	-----------------------	----------------------	-----------------------	-----------------------	-----------------------	-----------------------

							O	
			IIIA	IVA	VA	VIA	VIIA	He 2 4.00260
		10.81	12.011	14.0067	15.9994	18.998403	20.179	
		B 5	C 6	N 7	O 8	F 9	Ne 10	
		26.98154	28.0855	30.97376	32.06	35.453	39.948	
		Al 13	Si 14	P 15	S 16	Cl 17	Ar 18	
IB	IIB							
63.546	65.38	69.72	72.59	74.9216	78.96	79.904	83.80	
Cu 29	Zn 30	Ga 31	Ge 32	As 33	Se 34	Br 35	Kr 36	
107.868	112.41	114.82	118.69	121.75	127.60	126.9045	131.30	
Ag 47	Cd 48	In 49	Sn 50	Sb 51	Te 52	I 53	Xe 54	
196.9665	200.59	204.37	207.2	208.9804	208.98243	209.987	222.01761	
Au 79	Hg 80	Tl 81	Pb 82	Bi 83	Po 84	At 85	Rn 86	

167.26	168.9342	173.04	174.97
Er 68	Tm 69	Yb 70	Lu 71

257.09515	258	259	260
Fm 100	Md 101	No 102	Lr 103

Ac

Actinium

89

227.02777

IIIB

44 9592
Sc 21
88 9059
Y 39
138 9055
La 57
227 02777
Ac 89

Actinio

Actinium

Aktinium

Actinio

АКТИНИЙ

אקטיןיום

鋼
ア
ク
チ
ニ
ウム

Naturally occurring isotope: 227 (minute quantities only)

Density: 10.07 g/cm³ (25°C)

Melting point: 1100 ± 50°C **Boiling point:** 3200 ± 300°C (est)

Latent heat of fusion: 62 J/g

Specific heat: 0.12 J/g/°C

Thermal conductivity: 0.12 w/cm/°C (25°C)

Ionization potential (1st): 5.17 eV

Oxidation potential: Ac → Ac³⁺ + 3e = 2.2 V

Chemical valence: 3

Electrochemical equivalents: 2.82347 g/amp-hr

Ionic radius: 1.119 Å (Ac³⁺)

Valence electron potential (−eV): 38.60 (Ac³⁺)

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p⁶ 6d¹ 7s²

Valence electrons: 6d¹ 7s²

Crystal form: Cubic, face centered

Half life: 21.77 years

Cross section σ: 810 ± 20 barns

Al

Aluminum

13

26.98154

IIIA	
10 81	B
5	
26.98154	Al
13	
69 72	Ga
31	
114 82	In
49	
204 37	Tl
81	

Aluminio
Aluminium
Aluminium
Aluminio
алюминий

אלומין

鋁 アルミニウム

Naturally occurring isotope: 27

Density: 2.6984 g/cm³ (20°C)

Melting point: 660.37°C **Boiling point:** 2467°C

Latent heat of fusion: 395.7 J/g

Specific heat: 0.903 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 23.9 × 10⁻⁶cm/cm/°C (20°C)

Thermal conductivity: 2.37 w/cm/°C (25°C)

Electrical resistivity: 2.6548 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 5.986 eV

Electron work function ϕ : 4.28 eV

Oxidation potential: Al → Al³⁺ + 3e = 1.662 V

Chemical valence: 3

Electrochemical equivalents: 0.33556 g/amp-hr

Ionic radius: 0.535 Å (Al³⁺)

Valence electron potential (-eV): 80.7

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p¹

Valence electrons: 3s² 3p¹

Crystal form: Cubic, face centered

Cross section σ : 232 ± 3 mbarns

Vapor pressure: 2.42 × 10⁻⁶ Pa (at melting point)

Am

Americium

95

243.0614

Actinide Series

232.03807 Th 90	231.0358 Pa 91	238.029 U 92	237.0462 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08905 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Americio
Américium
Amerizium
Americio
америций
אמריציום

錒
アメリカシウム

Naturally occurring isotopes: None**Density:** 13.67 g/cm³ (20°C)**Melting point:** 1176°C **Boiling point:** 2011°C**Ionization potential (1st):** 5.99 eV**Oxidation potential:** Am → Am³⁺ + 3e = 2.32 V**Chemical valence:** 2, 3, 4, 5, 6**Electrochemical equivalents:** 3.0229 g/amp-hr**Ionic radius:** 0.982 Å (Am³⁺)**Valence electron potential (−eV):** 44.0**Principal quantum number:** 7**Principal electron shells:** K L M N O P Q**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f⁷ 6s² 6p⁶ 7s²**Valence electrons:** 5f⁷ 7s²**Crystal form:** Hexagonal**Half life:** 7.32 × 10³ years**Cross section σ:** 180 ± 20 barns

Sb

Antimony

51

121.75

VA

14.0067 N 7
30.97376 P 15
74.9216 As 33
121.75 Sb 51
208.9804 Bi 83

Antimônio

Antimoine

Antimon

Antimonio

сурьма

אנטיםן

銻
アンチモン

Naturally occurring isotopes: 121, 123

Density: 6.691 g/cm³ (20°C)

Melting point: 630.74°C **Boiling point:** 1750°C

Latent heat of fusion: 165.0 J/g

Specific heat: 0.207 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 9.2×10^{-6} cm/cm/°C (0°C)

Thermal conductivity: 0.244 w/cm/°C (25°C)

Electrical resistivity: 39×10^{-6} ohm-cm (0°C)

Ionization potential (1st): 8.641 eV

Electron work function ϕ : 4.55 eV

Oxidation potential: $2\text{Sb} + 3\text{H}_2\text{O} \rightarrow \text{Sb}_2\text{O}_3 + 6\text{H}^+ + 6\text{e}^- = -0.152 \text{ V}$

Chemical valence: -3, 0, 3, 5

Electrochemical equivalents: 1.5142 g/amp-hr

Ionic radius: 0.76 Å (Sb³⁺)

Valence electron potential (-eV): 57

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^3$

Valence electrons: $5s^2 5p^3$

Crystal form: Rhombohedral

Cross section σ : 5 ± 1 barns

Vapor pressure: 2.49×10^{-9} Pa (at melting point)

Ar

Argon

18

39.948

0
4 00260 He 2
20 179 Ne 10
39 948 Ar 18
83 80 Kr 36
131 30 Xe 54
222 01761 Rn 86

Argônio

Argon

Argon

Argón

аргон

ארגון

氬 アルゴン

Naturally occurring isotopes: 40, 36, 38

Density: 1.65 g/cm³ (-233°C), 1.784 × 10⁻³ g/cm³ (0°C)

Melting point: -189.2°C **Boiling point:** -185.7°C

Latent heat of fusion: 29.45 J/g

Specific heat: 0.52032 J/g°C (25°C)

Thermal conductivity: 0.1772 mw/cm°C (27°C at 1 atm)

Ionization potential (1st): 15.759 eV

Chemical valence: 0

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶

Valence electrons: (3s² 3p⁶)

Crystal form: Cubic, face centered

Cross section σ : 0.66 barns

As

Arsenic

33

74.9216

VA

14.0067 N 7
30.97376 P 15
74.9216 As 33
121.75 Sb 51
208.9804 Bi 83

Arsênio

Arsenic

Arsen

Arsénico

мышьяк

ᲐᲗᲗ

砷 砒素

Naturally occurring isotope: 75

Density: 5.73 g/cm³ (gray) (20°C)

Melting point: 817°C (at 28 atm) **Boiling point:** 613°C (sublimes)

Latent heat of fusion: 369.9 J/g

Specific heat: 0.329 J/g/°C (gray) (25°C)

Coefficient of lineal thermal expansion: 6.02 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.502 w/cm/°C (gray) (25°C)

Electrical resistivity: 35 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 9.81 eV

Electron work function ϕ : 3.75 eV

Oxidation potential: As + 2H₂O → HAsO₂ + 3H⁺ + 3e = -0.2476 V

Chemical valence: -3, 0, 3, 5

Electrochemical equivalents: 0.93177 g/amp-hr

Ionic radius: 0.58 Å (As³⁺)

Valence electron potential (-eV): 74

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p³

Valence electrons: 4s² 4p³

Crystal form: Rhombohedral

Cross section σ : 4.30 ± 0.10 barns

At

Astatine

85

209.987

VIIA

18 998403
F 9
35 453
Cl 17
79 904
Br 35
126 9045
I 53
209 987
At 85

Astato
Astatine
Astat
Astatino
АСТАТИН
АСТАТИН

砒
ア
ス
タ
チ
ン

Naturally occurring isotopes: None

Melting point: 302°C (est) **Boiling point:** 337°C (est)

Latent heat of fusion: 114 J/g (est)

Ionization potential (1st): 9.65 eV

Oxidation potential: $2\text{At}^- \rightarrow \text{At}_2 + 2e^- = -0.2 \text{ V}$

Chemical valence: 1, 3, 5, 7

Electrochemical equivalents: 7.8346 g/amp-hr

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 6s^2 6p^5$

Valence electrons: $6s^2 6p^5$

Half life: 8.1 hr

Ba

Barium

56

137.34

IIA

9 01218 Be 4
24 305 Mg 12
40 08 Ca 20
87 62 Sr 38
137 34 Ba 56
226-02544 Ra 88

Bárió

Barium

Barium

Bario

барий

בריום

鋇
バリウム

Naturally occurring isotopes: 138, 137, 136, 135, 134, 130, 132

Density: 3.59 g/cm³ (20°C)

Melting point: 725°C **Boiling point:** 1640°C

Latent heat of fusion: 55.79 J/g

Specific heat: 0.204 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 19.0 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.184 w/cm/°C (22°C)

Ionization potential (1st): 5.212 eV

Electron work function ϕ : 2.7 eV

Oxidation potential: Ba → Ba²⁺ + 2e = 2.906 V

Chemical valence: 2

Electrochemical equivalents: 2.5621 g/amp-hr

Ionic radius: 1.35 Å (Ba²⁺)

Valence electron potential (-εV): 21.3

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p⁶ 6s²

Valence electrons: 6s²

Crystal form: Cubic, body centered

Cross section σ : 1.2 ± 0.1 barns

Vapor pressure: 9.80 × 10 Pa (at melting point)

Bk

Berkelium

97

247.07032

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Berquélio

Berkelium

Berkelium

Berkelio

беркелий

ברקליום

錒
バーク
クリウ
ム

Naturally occurring isotopes: None

Density: 14.78 g/cm³ (25°C)

Melting point: 986 ± 25°C

Ionization potential (1st): 6.23 eV

Oxidation potential: Bk → Bk³⁺ + 3e = 1.97 V

Chemical valence: 3, 4

Electrochemical equivalents: 3.0727 g/amp-hr

Ionic radius: 0.949 Å (Bk³⁺)

Valence electron potential (-eV): 45.5

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f⁸ 6s² 6p⁶ 6d¹ 7s²

Valence electrons: 5f⁸ 6d¹ 7s²

Crystal form: Hexagonal

Half life: 1.4 × 10³ years

Be

Beryllium

4

9.01218

IIA

9.01218 Be 4
24.305 Mg 12
40.08 Ca 20
87.62 Sr 38
137.34 Ba 56
226.02544 Ra 88

Berilio

Beryllium

Beryllium

Berilio

бериллий

בריליום

鍍
バリ
リ
ウム

Naturally occurring isotope: 9

Density: 1.848 g/cm³ (20°C)

Melting point: 1278 ± 5°C **Boiling point:** 2970°C

Latent heat of fusion: 1301 J/g

Specific heat: 1.82 J/g°C (25°C)

Coefficient of lineal thermal expansion: 11.6 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 2.01 w/cm°C (25°C)

Electrical resistivity: 4.0 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 9.322 eV

Electron work function ϕ : 4.98 eV

Oxidation potential: Be → Be²⁺ + 2e = 1.85 V

Chemical valence: 2

Electrochemical equivalents: 0.16812 g/amp-hr

Ionic radius: 0.35 Å (Be²⁺)

Valence electron potential (-εV): 82

Principal quantum number: 2

Principal electron shells: K L

Electronic configuration: 1s² 2s²

Valence electrons: 2s²

Crystal form: Hexagonal, close packed

Cross section σ : 9.2 ± 0.5 mbarns

Vapor pressure: 4.18 Pa (at melting point)

Bi

Bismuth

83

208.9804

VA

14.0067
N
7
30.97376
P
15
74.9216
As
33
121.75
Sb
51
208.9804
Bi
83

Bismuto

Bismuth

Wismut

Bismuto

висмут

בִּיזְמוּט

鉍 ビスマス

Naturally occurring isotope: 209

Density: 9.78 g/cm³ (20°C)

Melting point: 271.3°C **Boiling point:** 1560 ± 5°C

Latent heat of fusion: 52.09 J/g

Specific heat: 0.122 J/g°C (25°C)

Coefficient of lineal thermal expansion: 13.3 × 10⁻⁶ cm/cm°C

Thermal conductivity: 0.0792 w/cm°C (25°C)

Electrical resistivity: 106.8 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 7.289 eV

Electron work function ϕ : 4.22 eV

Oxidation potential: Bi + H₂O → BiO⁺ + 2H⁺ + 3e⁻ = -0.320 V

Chemical valence: 3, 5

Electrochemical equivalents: 2.5990 g/amp-hr

Ionic radius: 1.03 Å (Bi³⁺)

Valence electron potential (-eV): 41.9

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p³

Valence electrons: 6s² 6p³

Crystal form: Rhombohedral

Cross section σ : 19 ± 2 mbarns

Vapor pressure: 6.27 × 10⁻⁴ Pa (at melting point)

B

Boron

5

10.81

IIIA	
10.81	B 5
26.98154	Al 13
69.72	Ga 31
114.82	In 49
204.37	Tl 81

Bóro

Bore

Bor

Boro

6op

בוֹר

硼素

Naturally occurring isotopes: 11, 10

Density: 2.34 g/cm³ (crystalline), 2.37 g/cm³ (amorphous) (both at 20°C)

Melting point: 2300°C **Boiling point:** 2550°C (sublimes)

Latent heat of fusion: 890.8 J/g

Specific heat: 1.03 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 8.3×10^{-6} cm/cm/°C (20°C)

Thermal conductivity: 0.274 w/cm/°C (25°C)

Electrical resistivity: 1.8×10^6 ohm-cm (0°C)

Ionization potential (1st): 8.298 eV

Electron work function ϕ : 4.45 eV

Oxidation potential: $B + 3H_2O \rightarrow H_3BO_3 + 3H^+ + 3e = -0.8698$ V

Chemical valence: 3

Electrochemical equivalents: 0.1344 g/amp-hr

Ionic radius: 0.23 Å (B³⁺)

Valence electron potential ($-\epsilon$ V): 190

Principal quantum number: 2

Principal electron shells: K L

Electronic configuration: $1s^2 2s^2 2p^1$

Valence electrons: $2s^2 2p^1$

Crystal form: Hexagonal, close packed

Cross section σ : 759 barns

Vapor pressure: 3.48×10^{-1} Pa (at melting point)

Br

Bromine

35

79.904

VIIA

18 996403
F 9
35 453
Cl 17
79 904
Br 35
126 9045
I 53
209 967
At 85

Bromo

Brome

Brom

Bromo

бром

ברום

溴 臭素

Naturally occurring isotopes: 79, 81

Density: 3.1028 g/cm³ (20°C)

Melting point: -7.2°C **Boiling point:** 58.78°C

Latent heat of fusion: 132.0 J/g (Br₂)

Specific heat: 0.47362 J/g/°C (Br₂) (25°C)

Thermal conductivity: 1.22 mw/cm/°C (27°C)

Electrical resistivity: 7.8 × 10¹² ohm-cm (0°C)

Ionization potential (1st): 11.814 eV

Oxidation potential: 2Br⁻ → Br₂ + 2e⁻ = -1.0652 V

Chemical valence: -1, 3, 5, 7

Electrochemical equivalents: 2.9812 g/amp-hr

Ionic radius: 1.96 Å (Br⁻)

Valence electron potential (-eV): -7.35

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁵

Valence electrons: 4s² 4p⁵

Crystal form: Orthorhombic, rhombic

Cross section σ: 6.8 ± 0.1 barns

Vapor pressure: 5.80 × 10³ Pa (at melting point)

Cd

Cadmium

48

112.41

HB

65 38 Zn 30
112.41 Cd 48
200 59 Hg 80

Cádmio

Cadmium

Cadmium

Cadmio

кадмий

קדמיום

カ
ド
ミ
ウ
ム

Naturally occurring isotopes: 114, 112, 111, 110, 113, 116, 106, 108

Density: 8.65 g/cm³ (20°C)

Melting point: 320.9°C **Boiling point:** 765°C

Latent heat of fusion: 54.01 J/g

Specific heat: 0.231 J/g°C (25°C)

Coefficient of lineal thermal expansion: 29.8 × 10⁻⁶ cm/cm°C (25°C)

Thermal conductivity: 0.969 w/cm°C (25°C)

Electrical resistivity: 6.83 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 8.993 eV

Electron work function ϕ : 4.22 eV

Oxidation potential: Cd → Cd²⁺ + 2e = 0.4029 V

Chemical valence: 2

Electrochemical equivalents: 2.0970 g/amp-hr

Ionic radius: 0.97 Å (Cd²⁺)

Valence electron potential (-εV): 30

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s²

Valence electrons: 5s²

Crystal form: Hexagonal, close packed

Cross section σ : 2450 ± 20 barns

Vapor pressure: 1.48 × 10 Pa (at melting point)

Ca

Calcium

20

40.08

IIA

9 01218 Be 4
24 305 Mg 12
40 08 Ca 20
87 62 Sr 38
137 34 Ba 56
226.07544 Ra 88

Cálcio
Calcium
Kalzium
Calcio
кальций

ἵπῶ

鈣 鈣
カルシウム

Naturally occurring isotopes: 40, 44, 42, 48, 43, 46

Density: 1.55 g/cm³ (20°C)

Melting point: 839 ± 2°C **Boiling point:** 1484°C

Latent heat of fusion: 216.2 J/g

Specific heat: 0.632 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 22.3 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 2.01 w/cm/°C (25°C)

Electrical resistivity: 3.91 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 6.113 eV

Electron work function ϕ : 2.87 eV

Oxidation potential: Ca → Ca²⁺ + 2e = 2.866 V

Chemical valence: 2

Electrochemical equivalents: 0.7477 g/amp-hr

Ionic radius: 0.99 Å (Ca²⁺)

Valence electron potential (-eV): 29

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 4s²

Valence electrons: 4s²

Crystal form: Cubic, face centered

Cross section σ : 0.44 ± 0.02 barns

Vapor pressure: 2.54 × 10² Pa (at melting point)

Cf

Californium

98

251.07961

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Californio
Californium
Californium
Californio
калифорний
קליפורניום

金開

カリ
フォ
リニ
ウム

Naturally occurring isotopes: None

Density: 15.1 g/cm³ (25°C)

Melting point: 900 ± 30°C

Ionization potential (1st): 6.30 eV

Oxidation potential: Cf → Cf³⁺ + 3e = 2.0 V

Chemical valence: 2, 3, 4

Electrochemical equivalents: 3.1226 g/amp-hr

Ionic radius: 0.934 Å (Cf³⁺)

Valence electron potential (-εV): 44.5

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f¹⁰ 6s² 6p⁶ 7s²

Valence electrons: 5f¹⁰ 7s²

Crystal form: Hexagonal

Half life: 900 years

Cross section σ: 2100 ± 1000 barns

C

Carbon

6

12.011

IVA

12 011 C 6
28 0855 Si 14
72 59 Ge 32
118 69 Sn 50
207.2 Pb 82

Carbono

Carbone

Kohlenstoff

Carbono

углерод

꠆꠆꠆

碳	炭
素	素

Naturally occurring isotopes: 12, 13, 14**Density:** 3.52 g/cm³ (diamond), 1.9–2.3 g/cm³ (graphite), 1.8–2.1 g/cm³ (amorphous) (all at 20°C)**Melting point:** 3550°C **Boiling point:** 4827°C**Specific heat:** 0.7099 J/g°C (graphite) (25°C)**Coefficient of lineal thermal expansion:** 2.10 × 10⁻⁶ cm/cm°C (graphite) (30°C)**Thermal conductivity:** 0.8–2.2 w/cm°C (graphite) (25°C)**Electrical resistivity:** 1375 × 10⁻⁶ ohm-cm (graphite) (0°C)**Ionization potential (1st):** 11.260 eV**Electron work function ϕ :** 5.0 eV**Oxidation potential:** CH₄ → C + 4H⁺ + 4e⁻ = -0.1316 V**Chemical valence:** 2, 3, 4**Electrochemical equivalents:** 0.11203 g/amp-hr**Ionic radius:** 0.16 Å (C⁴⁺)**Valence electron potential (-eV):** 360**Principal quantum number:** 2**Principal electron shells:** K L**Electronic configuration:** 1s² 2s² 2p²**Valence electrons:** 2s² 2p²**Crystal form:** Hexagonal (graphite), cubic (diamond)**Cross section σ :** 3.4 ± 0.2 mbarns

Ce

Cerium

58

140.12

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Cério

Cérium

Zerium

Cerio

церий

צריום

 銻 セリ
ウ
ム
Naturally occurring isotopes: 140, 142, 138, 136**Density:** 6.657 g/cm³ (25°C)**Melting point:** 799°C **Boiling point:** 3426°C**Latent heat of fusion:** 65.7 J/g**Specific heat:** 0.192 J/g/°C (25°C)**Coefficient of linear thermal expansion:** 7.1 × 10⁻⁶ cm/cm/°C (25°C)**Thermal conductivity:** 0.113 w/cm/°C (25°C)**Electrical resistivity:** 77 × 10⁻⁶ ohm-cm (25°C)**Ionization potential (1st):** 5.47 eV**Electron work function ϕ :** 2.84 eV**Oxidation potential:** Ce → Ce³⁺ + 3e = 2.483 V**Chemical valence:** 3, 4**Electrochemical equivalents:** 1.7426 g/amp-hr**Ionic radius:** 1.034 Å (Ce³⁺)**Valence electron potential (-εV):** 41.78**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f² 5s² 5p⁶ 6s²**Valence electrons:** 4f² 6s²**Crystal form:** Cubic, face centered**Cross section σ :** 0.73 ± 0.08 barns

Cs

Cesium

55

132.9054

IA

1 0079 H 1
6 941 Li 3
22 98977 Na 11
39 098 K 19
85 4678 Rb 37
132 9054 Cs 55
223 019/6 Fr 87

Césio

Césium

Caesium

Ceslo

цезий

סזיום

 銫
セシウム
Naturally occurring isotope: 133**Density:** 1.873 g/cm³ (20°C)**Melting point:** 28.40 ± 0.01°C **Boiling point:** 669.3°C**Latent heat of fusion:** 16.372 J/g**Specific heat:** 0.241 J/g/°C (25°C)**Coefficient of lineal thermal expansion:** 97 × 10⁻⁶ cm/cm/°C (20°C)**Thermal conductivity:** 0.359 w/cm/°C (solid at melting point)**Electrical resistivity:** 20.46 × 10⁻⁶ ohm-cm (20°C)**Ionization potential (1st):** 3.894 eV**Electron work function ϕ :** 2.14 eV**Oxidation potential:** Cs → Cs⁺ + e = 2.923 V**Chemical valence:** 1**Electrochemical equivalents:** 4.95870 g/amp-hr**Ionic radius:** 1.67 Å (Cs⁺)**Valence electron potential (-εV):** 8.62**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p⁶ 6s¹**Valence electrons:** 6s¹**Crystal form:** Cubic, body centered**Cross section σ :** 30.0 ± 1.5 barns**Vapor pressure:** 2.50 × 10⁻⁵ Pa (at melting point)

Cl

Chlorine

17

35.453

VIIA

18 998403
F 9
35 453
Cl 17
79 904
Br 35
126 9045
I 53
209 987
At 85

Clóro

Chlore

Chlor

Cloro

חסר

כלור

氯 塩素

Naturally occurring isotopes: 35, 37

Density: 1.56 g/cm³ (−33.6°C), 3.214 × 10^{−3} g/cm³ (0°C)

Melting point: −100.98°C **Boiling point:** −34.6°C

Latent heat of fusion: 180.8 J/g (Cl₂)

Specific heat: 0.4782 J/g/°C (Cl₂) (25°C)

Thermal conductivity: 0.089 mw/cm/°C (27°C at 1 atm)

Ionization potential (1st): 12.967 eV

Oxidation potential: 2Cl[−] → Cl₂ + 2e[−] = −1.3595 V

Chemical valence: −1, 3, 5, 7

Electrochemical equivalents: 1.3228 g/amp-hr

Ionic radius: 1.81 Å (Cl[−])

Valence electron potential (−εV): −7.96

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁵

Valence electrons: 3s² 3p⁵

Crystal form: Tetragonal

Cross section σ: 33 barns

Vapor pressure: 1.30 × 10³ Pa (at melting point)

Cr

Chromium

24

51.996

VIB

51.996
Cr 24
95.94
Mo 42
183.85
W 74
106

Crômio

Chrom

Chrom

Cromo

хром

כרום

鉻
ウ
ロ
ー
ム

Naturally occurring isotopes: 52, 53, 50, 54

Density: 7.20 g/cm³ (20°C)

Melting point: 1857 ± 20°C **Boiling point:** 2672°C

Latent heat of fusion: 265.7 J/g

Specific heat: 0.449 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 6.2 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.939 w/cm/°C (25°C)

Electrical resistivity: 12.9 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 6.766 eV

Electron work function ϕ : 4.5 eV

Oxidation potential: Cr → Cr³⁺ + 3e = 0.744 V

Chemical valence: 1, 2, 3, 4, 5, 6

Electrochemical equivalents: 0.32333 g/amp-hr

Ionic radius: 0.52 Å (Cr⁶⁺)

Valence electron potential (-eV): 170

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁵ 4s¹

Valence electrons: 3d⁵ 4s¹

Crystal form: Cubic, body centered

Cross section σ : 3.1 ± 0.2 barns

Vapor pressure: 9.90 × 10² Pa (at melting point)

Co

Cobalt

27

58.9332

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
	109	

Cobalto

Cobalt

Kobalt

Cobalto

кобальт

קובלט

钴
コバルト

Naturally occurring isotope: 59

Density: 8.71 g/cm³ (21°C)

Melting point: 1495°C **Boiling point:** 2870°C

Latent heat of fusion: 258.6 J/g

Specific heat: 4.21 J/g°C (25°C)

Coefficient of lineal thermal expansion: 13.80 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 1.00 w/cm°C (25°C)

Electrical resistivity: 6.24 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.86 eV

Electron work function ϕ : 5.0 eV

Oxidation potential: Co → Co²⁺ + 2e = 0.277 V

Chemical valence: 2, 3, 4

Electrochemical equivalents: 1.0994 g/amp-hr

Ionic radius: 0.745 Å (Co³⁺)

Valence electron potential (-eV): 38.7

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁷ 4s²

Valence electrons: 3d⁷ 4s²

Crystal form: Hexagonal, close packed

Cross section σ : 37.5 ± 0.2 barns

Vapor pressure: 1.75 × 10² Pa (at melting point)

Cu

Copper

29

63.546

IB

63 546
Cu
29
107 868
Ag
47
196 9665
Au
79

Cobre

Cuivre

Kupfer

Cobre

медь

נחשת

銅 銅

Naturally occurring isotopes: 63, 65

Density: 8.96 g/cm³ (25°C)

Melting point: 1083.4 ± 0.2°C **Boiling point:** 2567°C

Latent heat of fusion: 205.6 J/g

Specific heat: 0.3845 J/g°C (25°C)

Coefficient of lineal thermal expansion: 16.6 × 10⁻⁶ cm/cm°C (25°C)

Thermal conductivity: 4.01 w/cm°C (25°C)

Electrical resistivity: 1.678 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.726 eV

Electron work function ϕ : 4.65 eV

Oxidation potentials: Cu → Cu⁺ + ϵ = -0.521 V

Cu → Cu²⁺ + 2 ϵ = -0.3419 V

Chemical valence: 1, 2

Electrochemical equivalents: 1.1855 g/amp-hr

Ionic radius: 0.73 Å (Cu²⁺)

Valence electron potential (-eV): 34

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s¹

Valence electrons: 3d¹⁰ 4s¹

Crystal form: Cubic, face centered

Cross section σ : 3.8 ± 0.1 barns

Vapor pressure: 5.05 × 10⁻² Pa (at melting point)

Cm

Curium

96

247.07038

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.0289 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Cúrio

Curium

Curium

Curio

кюриий

קיוריום

銅
キュ
リ
ウム

Naturally occurring isotopes: None

Density: 13.51 g/cm³ (25°C)

Melting point: 1340 ± 40°C **Boiling point:** 3110°C

Ionization potential (1st): 6.02 eV

Oxidation potential: Cm → Cm³⁺ + 3e⁻ = 2.07 V

Chemical valence: 3, 4

Electrochemical equivalents: 3.0727 g/amp-hr

Ionic radius: 0.970 Å (Cm³⁺)

Valence electron potential (-εV): 44.5

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f⁷ 6s² 6p⁶ 6d¹ 7s²

Valence electrons: 5f⁷ 6d¹ 7s²

Crystal form: Hexagonal

Half life: 1.6 × 10⁷ years

Cross section σ: 180 barns

Dy

Dysprosium

66

162.50

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Disprósio
Dysprosium
Dysprosium
Disprosio
диспрозий
דיטברויזיום

鎢
ジ
ス
プ
ロ
シ
ウ
ム

Naturally occurring isotopes: 164, 162, 163, 161, 160, 158, 156

Density: 8.550 g/cm³ (25°C)

Melting point: 1412°C **Boiling point:** 2562°C

Latent heat of fusion: 105.6 J/g

Specific heat: 173 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 8.6×10^{-6} cm/cm/°C (25°C)

Thermal conductivity: 0.107 w/cm/°C (25°C)

Electrical resistivity: 90×10^{-6} ohm-cm (25°C)

Ionization potential (1st): 5.928 eV

Oxidation potential: Dy → Dy³⁺ + 3e = 2.353 V

Chemical valence: 3

Electrochemical equivalents: 2.0210 g/amp-hr

Ionic radius: 0.912 Å (Dy³⁺)

Valence electron potential (-eV): 47.4

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁰ 5s²
5p⁶ 6s²

Valence electrons: 4f¹⁰ 6s²

Crystal form: Hexagonal, close packed

Cross section σ : 930 ± 20 barns

Es

Einsteinium

99

254.08805

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Einstênio
Einsteinium
Einsteinium
Einsteinio
ЭЙНШТЕЙНИЙ
אֵינשטייניום

𐤀
𐤅
𐤍
𐤏
𐤑
𐤓
𐤕
𐤗
𐤙
𐤛
𐤝
𐤟
𐤡
𐤣
𐤥
𐤨
𐤩
𐤫
𐤬
𐤭
𐤯
𐤱
𐤳
𐤵
𐤷
𐤹
𐤻
𐤽
𐤿

Naturally occurring isotopes: None

Melting point: $860 \pm 30^\circ\text{C}$

Ionization potential (1st): 6.42 eV

Oxidation potential: $\text{Es} \rightarrow \text{Es}^{2+} + 2e = 2.3 \text{ V}$

Chemical valence: 2, 3

Electrochemical equivalents: 4.7400 g/amp-hr

Ionic radius: 0.925 Å (Es^{3+})

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 5f^{11} 6s^2 6p^6 7s^2$

Valence electrons: $5f^{11} 7s^2$

Crystal form: Cubic, face centered

Half life: 276 days

Cross section σ : < 40 barns

Er

Erbium

68

167.26

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Érbio

Erbium

Erbium

Erbio

эрбий

ארביום

鉬 エルビウム

Naturally occurring isotopes: 166, 168, 167, 170, 164, 162

Density: 9.066 g/cm³ (25°C)

Melting point: 1529°C **Boiling point:** 2863°C

Latent heat of fusion: 102.6 J/g

Specific heat: 0.168 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 9.2×10^{-6} cm/cm/°C (25°C)

Thermal conductivity: 0.145 w/cm/°C (25°C)

Electrical resistivity: 107.0×10^{-6} ohm-cm (25°C)

Ionization potential (1st): 6.10 eV

Oxidation potential: $Er \rightarrow Er^{3+} + 3e = 2.296$ V

Chemical valence: 3

Electrochemical equivalents: 2.0802 g/amp-hr

Ionic radius: 0.881 Å (Er³⁺)

Valence electron potential (-eV): 49.0

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{12} 5s^2 5p^6 6s^2$

Valence electrons: $4f^{12} 6s^2$

Crystal form: Hexagonal, close packed

Cross section σ : 160 ± 30 barns

Eu

Europium

63

151.96

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Európio
Europium
Europium
Europio
европий
אירופיום

鎔 ユーロピウム

Naturally occurring isotopes: 153, 151

Density: 5.243 g/cm³ (25°C)

Melting point: 822°C **Boiling point:** 1597°C

Latent heat of fusion: 68.9 J/g

Specific heat: 0.182 J/g°C (25°C)

Coefficient of lineal thermal expansion: 26 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 0.139 w/cm°C (25°C)

Electrical resistivity: 81 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 5.666 eV

Electron work function φ: 2.5 eV

Oxidation potential: Eu → Eu³⁺ + 3e = 2.407 V

Chemical valence: 2, 3

Electrochemical equivalents: 1.8899 g/amp-hr

Ionic radius: 0.947 Å (Eu³⁺)

Valence electron potential (-εV): 45.6

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f⁷ 5s² 5p⁶ 6s²

Valence electrons: 4f⁷ 6s²

Crystal form: Cubic, body centered

Cross section σ: 4100 ± 100 barns

Vapor pressure: 1.44 × 10² Pa (at melting point)

Fm

Fermium

100

257.09515

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Férmio

Fermium

Fermium

Fermio

фермий

פרמיום

鍳

フェリミウム

Naturally occurring isotopes: None**Ionization potential (1st):** 6.50 eV**Oxidation potential:** $\text{Fm} \rightarrow \text{Fm}^{3+} + 3\epsilon = 2.0 \text{ V}$ **Chemical valence:** 2, 3**Electrochemical equivalents:** 3.1974 g/amp-hr**Principal quantum number:** 7**Principal electron shells:** K L M N O P Q**Electronic configuration:** $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 5f^{12} 6s^2 6p^6 7s^2$ **Valence electrons:** $5f^{12} 7s^2$ **Half life:** 80 days

F Fluorine

9

18.998403

VIIA

18 998403
F 9
35 453
Cl 17
79 904
Br 35
126 9045
I 53
209 987
At 85

Flúor

Fluor

Fluor

Flúor

φτορ

פלוואר

氟 弗素

Naturally occurring isotope: 19

Density: $1.696 \times 10^{-3} \text{ g/cm}^3$ (0°C)

Melting point: -219.62°C **Boiling point:** -188.14°C

Latent heat of fusion: 26.89 J/g (F_2)

Specific heat: 0.824 J/g/°C (F_2) (25°C)

Thermal conductivity: 0.279 mW/cm/°C (27°C at 1 atm)

Ionization potential (1st): 17.422 eV

Oxidation potential: $\text{F}^- \rightarrow \frac{1}{2}\text{F}_2 + \epsilon = -2.87 \text{ V}$

Chemical valence: - 1

Electrochemical equivalents: 0.70883 g/amp-hr

Ionic radius: 1.33 Å (F^-)

Valence electron potential ($-\epsilon\text{V}$): - 10.1

Principal quantum number: 2

Principal electron shells: K L

Electronic configuration: $1s^2 2s^2 2p^5$

Valence electrons: $2s^2 2p^5$

Cross section σ : 9.8 ± 0.7 mbarns

Vapor pressure: $4.90 \times 10^2 \text{ Pa}$ (at melting point)

Fr

Francium

87

223.01976

IA

1 0079	H 1
6 941	Li 3
22 98977	Na 11
39 098	K 19
85 4678	Rb 37
132 9054	Cs 55
223 01976	Fr 87

Francio
 Francium
 Franzium
 Francio
 франций
 פרנציום
 釷
 ワ
 ラ
 ン
 シ
 ウ
 ム

Naturally occurring isotopes: None (actinium decay product)

Melting point: 27°C (est) **Boiling point:** 677°C (est)

Latent heat of fusion: 9.39 J/g (est)

Ionization potential (1st): 3.83 eV

Chemical valence: 1

Electrochemical equivalents: 8.3209 g/amp-hr

Ionic radius: 1.80 Å (Fr⁺)

Valence electron potential (−εV): 8.00

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
 5d¹⁰ 6s² 6p⁶ 7s¹

Valence electrons: (7s¹)

Half life: 22 minutes

Crystal form: Cubic, body centered

Gd

Gadolinium

64

157.25

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Gadolínio
Gadolinium
Gadolinium
Gadolinio
гадолиний
גדוליניום

釷
ガ
ド
リ
ニ
ウ
ム

Naturally occurring isotopes: 158, 160, 156, 157, 155, 154, 152

Density: 7.900 g/cm³ (25°C)

Melting point: 1313°C **Boiling point:** 3266°C

Latent heat of fusion: 98.51 J/g

Specific heat: 0.235 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 9.7 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.105 w/cm/°C (25°C)

Electrical resistivity: 140.5 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.14 eV

Electron work function φ: 3.1 eV

Oxidation potential: Gd → Gd³⁺ + 3e = 2.397 V

Chemical valence: 3

Electrochemical equivalents: 1.9557 g/amp-hr

Ionic radius: 0.938 Å (Gd³⁺)

Valence electron potential (-eV): 46.1

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f⁷ 5s² 5p⁶ 5d¹ 6s²

Valence electrons: 4f⁷ 5d¹ 6s²

Crystal form: Hexagonal, close packed

Cross section σ: 46,000 ± 2000 barns

Vapor pressure: 2.44 × 10⁴ Pa (at melting point)

Ga

Gallium

31

69.72

IIIA	
10 81 B 5	
26 98 154 Al 13	
69 72 Ga 31	
114 82 In 49	
204 37 Tl 81	

Gálio
Gallium
Gallium
Galio
галлий

גליום

鎳 ガリウム

Naturally occurring isotopes: 69, 71

Density: 5.906 g/cm³ (25°C)

Melting point: 29.78°C **Boiling point:** 2403°C

Latent heat of fusion: 80.17 J/g

Specific heat: 0.371 J/g°C (25°C)

Coefficient of lineal thermal expansion: 18.1×10^{-6} cm/cm°C (25°C)

Thermal conductivity: 0.281 w/cm°C (liquid) (30°C)

Electrical resistivity: 17.4×10^{-6} ohm-cm (20°C)

Ionization potential (1st): 5.999 eV

Electron work function ϕ : 4.2 eV

Oxidation potential: $\text{Ga} \rightarrow \text{Ga}^{3+} + 3\text{e} = -0.529 \text{ V}$

Chemical valence: 2, 3

Electrochemical equivalents: 0.8671 g/amp-hr

Ionic radius: 0.620 Å (Ga^{3+})

Valence electron potential ($-\epsilon\text{V}$): 69.7

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^1$

Valence electrons: $4s^2 4p^1$

Crystal form: Orthorhombic, rhombic

Cross section σ : 3.1 ± 0.3 barns

Vapor pressure: 9.31×10^{-36} Pa (at melting point)

Ge

Germanium

32

72.59

IVA

12 011 C 6
28 0855 Si 14
72 59 Ge 32
118 69 Sn 50
207 2 Pb 82

Germanio

Germanium

Germanium

Germanio

германий

גרמניום

 鍺
 ゲル
 マニ
 ウム
Naturally occurring isotopes: 74, 72, 70, 73, 76**Density:** 5.323 g/cm³ (25°C)**Melting point:** 937.4°C **Boiling point:** 2830°C**Latent heat of fusion:** 438.3 J/g**Specific heat:** 0.3216 J/g/°C (25°C)**Coefficient of lineal thermal expansion:** 5.75 × 10⁻⁶ cm/cm/°C (20°C)**Thermal conductivity:** 0.667 w/cm/°C (25°C)**Electrical resistivity:** 47 ohm-cm (intrinsic resistivity) (22°C)**Ionization potential (1st):** 7.899 eV**Electron work function ϕ :** 5.0 eV**Oxidation potential:** Ge + 2H₂O → GeO₂ + 4H⁺ + 4e⁻ = -0.15 V**Chemical valence:** -4, 2, 4**Electrochemical equivalents:** 0.6771 g/amp-hr**Ionic radius:** 0.530 Å (Ge⁴⁺)**Valence electron potential (-eV):** 109**Principal quantum number:** 4**Principal electron shells:** K L M N**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p²**Valence electrons:** 4s² 4p²**Crystal form:** Cubic, diamond**Cross section σ :** 2.30 ± 0.26 barns**Vapor pressure:** 7.46 × 10⁻⁵ Pa (at melting point)

Au

Gold

79

196.9665

IB
63.546 Cu 29
107.868 Ag 47
196.9665 Au 79

Ouro
Or
Gold
Oro
ЗОЛОТО
ግዓዝ

金 金

Naturally occurring isotope: 197

Density: 19.32 g/cm³ (20°C)

Melting point: 1064.43°C **Boiling point:** 3080°C

Latent heat of fusion: 62.81 J/g

Specific heat: 0.1290 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 14.2 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 3.19 w/cm/°C (25°C)

Electrical resistivity: 2.44 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 9.225 eV

Electron work function φ: 5.1 eV

Oxidation potentials: Au → Au⁺ + e = -1.691 V

Au → Au³⁺ + 3e = -1.498 V

Chemical valence: 1, 3

Electrochemical equivalents: 2.4496 g/amp-hr

Ionic radius: 0.85 Å (Au³⁺)

Valence electron potential (-eV): 51

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s¹

Valence electrons: 5d¹⁰ 6s¹

Crystal form: Cubic, face centered

Cross section σ: 98.8 ± 0.3 barns

Vapor pressure: 2.37 × 10⁻⁴ Pa (at melting point)

Hf

Hafnium

72

178.49

IVB

47.90 Ti 22
91.22 Zr 40
178.49 Hf 72
104

Háfnió

Hafnium

Hafnium

Hafnio

гафний

ハフニウム

鉛
ハ
フ
ニ
ウ
ム

Naturally occurring isotopes: 180, 178, 177, 179, 176, 174

Density: 13.31 g/cm³ (20°C)

Melting point: 2227 ± 20°C **Boiling point:** 4602°C

Latent heat of fusion: 122.0 J/g

Specific heat: 0.144 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 5.6 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.230 w/cm/°C (25°C)

Electrical resistivity: 35.1 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.65 eV

Electron work function φ: 3.9 eV

Oxidation potential: Hf → Hf⁴⁺ + 4e = 1.70 V

Chemical valence: 4

Electrochemical equivalents: 1.6649 g/amp-hr

Ionic radius: 0.71 Å (Hf⁴⁺)

Valence electron potential (-εV): 81

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d² 6s²

Valence electrons: 5d² 6s²

Crystal form: Hexagonal, close packed

Cross section σ: 103 ± 3 barns

Vapor pressure: 1.12 × 10⁻³ Pa (at melting point)

He

Helium

2

4.00260

0
4.00260 He 2
20.179 Ne 10
39.948 Ar 18
83.80 Kr 36
131.30 Xe 54
222.01761 Rn 86

Hélio

Hélium

Helium

Helio

гелий

הליום

氦
ヘリウム

Naturally occurring isotopes: 4, 3

Density: $0.17847 \times 10^{-3} \text{ g/cm}^3$ (0°C)

Melting point: -272.2°C (26 atm); | **Boiling point:** -268.934°C

Latent heat of fusion: 5.23 J/g

Specific heat: 5.1930 J/g°C (25°C)

Thermal conductivity: 1.520 mw/cm°C (25°C at 1 atm)

Ionization potential (1st): 24.58 eV

Chemical valence: 0

Principal quantum number: 1

Principal electron shells: K

Electronic configuration: $1s^2$

Valence electrons: ($1s^2$)

Crystal form: Hexagonal, close packed

Cross section σ : 0.007 barns

Ho

Holmium

67

164.9304

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Hólmio
Holmium
Holmium
Holmio
ХОЛМИЙ

הוֹלְמִיּוֹם

釹
ホルミウム

Naturally occurring isotope: 165

Density: 8.795 g/cm³ (25°C)

Melting point: 1474°C **Boiling point:** 2695°C

Latent heat of fusion: 104.1 J/g

Specific heat: 0.165 J/g°C (25°C)

Coefficient of lineal thermal expansion: 9.5 × 10⁻⁶ cm/cm°C (400°C)

Thermal conductivity: 0.162 w/cm°C (25°C)

Electrical resistivity: 87.0 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.02 eV

Oxidation potential: Ho → Ho³⁺ + 3e = 2.319 V

Chemical valence: 3

Electrochemical equivalents: 2.0512 g/amp-hr

Ionic radius: 0.901 Å (Ho³⁺)

Valence electron potential (-εV): 47.9

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹¹ 5s² 5p⁶ 6s²

Valence electrons: 4f¹¹ 6s²

Crystal form: Hexagonal, close packed

Cross section σ: 65 ± 2 barns

H

Hydrogen

1

1.0079

IA	
1.0079	H 1
6.941	Li 3
22.98977	Na 11
39.098	K 19
85.4678	Rb 37
132.9054	Cs 55
223.01976	Fr 87

Hidrogênio

Hydrogène

Wasserstoff

Hidrógeno

водород

מימן

氢气 水素

Naturally occurring isotopes: 1.007825 (protium), 2.01410 (deuterium), 3.01605 (tritium)

Density: $0.08988 \times 10^{-3} \text{ g/cm}^3$ (0°C)

Melting point: -259.14°C **Boiling point:** -252.87°C

Latent heat of fusion: 116.3 J/g (H_2)

Specific heat: 14.30 J/g/°C (H_2) (25°C)

Thermal conductivity: 1.815 mw/cm/°C (27°C at 1 atm)

Ionization potential (1st): 13.59765 eV

Oxidation potentials: $\text{H}_2 \rightarrow 2\text{H}^+ + \epsilon = 0.00000 \text{ V}$

$\text{H}^- \rightarrow \frac{1}{2}\text{H}_2 + \epsilon = 2.25 \text{ V}$

Chemical valence: 1

Electrochemical equivalents: 0.037605 g/amp-hr

Ionic radius: 0.012 Å (H^+)

Valence electron potential ($-\epsilon\text{V}$): 1200

Principal quantum number: 1

Principal electron shells: K

Electronic configuration: $1s^1$

Valence electrons: $1s^1$

Crystal form: Hexagonal, close packed

Cross section σ : 0.33 barns

In

Indium

49

114.82

IIIA	
10.81	B
5	
26.98154	Al
13	
69.72	Ga
31	
114.82	In
49	
204.37	Tl
81	

Índio

Indium

Indium

Indio

индий

இந்திய

 錫
 イ
 ソ
 ジ
 ウ
 ム
Naturally occurring isotopes: 115, 113**Density:** 7.28 g/cm³ (20°C)**Melting point:** 156.61°C **Boiling point:** 2080°C**Latent heat of fusion:** 28.44 J/g**Specific heat:** 0.233 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 24.8 × 10⁻⁶ cm/cm°C (20°C)**Thermal conductivity:** 0.818 w/cm°C (25°C)**Electrical resistivity:** 8.37 × 10⁻⁶ ohm-cm (0°C)**Ionization potential (1st):** 5.786 eV**Electron work function ϕ :** 4.12 eV**Oxidation potential:** In → In³⁺ + 3e = 0.343 V**Chemical valence:** 1, 2, 3**Electrochemical equivalents:** 1.4280 g/amp-hr**Ionic radius:** 0.800 Å (In³⁺)**Valence electron potential (-εV):** 54.0**Principal quantum number:** 5**Principal electron shells:** K L M N O**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p¹**Valence electrons:** 5s² 5p¹**Crystal form:** Tetragonal**Cross section σ :** 194 ± 2 barns**Vapor pressure:** 1.42 × 10¹⁷ Pa (at melting point)



Iodine

53

126.9045

VIIA

18 998403
F
9
35 453
Cl
17
79 904
Br
35
126 9045
I
53
209 987
At
85

Iodo

Iode

Iod

Yodo

iod

יוד

碘 沃素

Naturally occurring isotope: 127

Density: 4.93 g/cm³ (20°C)

Melting point: 113.5°C **Boiling point:** 184.35°C

Latent heat of fusion: 124.4 J/g (I₂)

Specific heat: 0.21448 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 93 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 4.49 mw/cm/°C (27°C)

Electrical resistivity: 1.3 × 10⁹ ohm-cm (20°C)

Ionization potential (1st): 10.451 eV

Oxidation potential: I⁻ → ½I₂ + e = -0.5355 V

Chemical valence: -1, 3, 5, 7

Electrochemical equivalents: 4.7348 g/amp-hr

Ionic radius: 2.20 Å (I⁻)

Valence electron potential (-eV): -6.55

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p⁵

Valence electrons: 5s² 5p⁵

Crystal form: Orthorhombic

Cross section σ: 6.2 ± 0.2 barns

Ir

Iridium

77

192.22

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
109		

Iridio

Iridium

Iridium

Iridio

ИРИДИЙ

אִירִידיּוּם

 銱
 イ
 リ
 ジ
 ム
Naturally occurring isotopes: 193, 191**Density:** 22.42 g/cm³ (17°C)**Melting point:** 2410°C **Boiling point:** 4130°C**Latent heat of fusion:** 137.2 J/g**Specific heat:** 0.131 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 6.6 × 10⁻⁶ cm/cm°C (25°C)**Thermal conductivity:** 1.47 w/cm°C (25°C)**Electrical resistivity:** 4.71 × 10⁻⁶ ohm-cm (20°C)**Ionization potential (1st):** 9.1 eV**Electron work function ϕ :** 5.27 eV**Oxidation potential:** Ir + 6Cl⁻ → IrCl₆³⁻ + 3e = -0.77 V**Chemical valence:** 2, 3, 4, 6**Electrochemical equivalents:** 1.793 g/amp-hr**Ionic radius:** 0.625 Å (Ir⁴⁺)**Valence electron potential (-eV):** 92.2**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d⁷ 6s²**Valence electrons:** 5d⁷ 6s²**Crystal form:** Cubic, face centered**Cross section σ :** 425 ± 15 barns**Vapor pressure:** 1.47 Pa (at melting point)

Fe

Iron

26

55.847

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
109		

Ferro

Fer

Eisen

Hierro

железо

ברזל

鐵 鐵

Naturally occurring isotopes: 56, 54, 57, 58

Density: 7.874 g/cm³ (20°C)

Melting point: 1535°C **Boiling point:** 2750°C

Latent heat of fusion: 275.1 J/g

Specific heat: 0.450 J/g°C (25°C)

Coefficient of lineal thermal expansion: 11.76 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.804 w/cm/°C (25°C)

Electrical resistivity: 9.71 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.870 eV

Electron work function ϕ : 4.70 eV

Oxidation potential: Fe → Fe²⁺ + 2e = 0.4402 V

Chemical valence: 2, 3, 4, 6

Electrochemical equivalents: 0.69455 g/amp-hr

Ionic radius: 0.645 Å (Fe³⁺)

Valence electron potential (-eV): 67.0

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁶ 4s²

Valence electrons: 3d⁶ 4s²

Crystal form: Cubic, body centered

Cross section σ : 2.56 ± 0.05 barns

Vapor pressure: 7.05 Pa (at melting point)

Kr

Krypton

36

83.80

O	
4.00260	He
2	
20.179	Ne
10	
39.948	Ar
18	
83.80	Kr
36	
131.30	Xe
54	
222.01761	Rn
86	

Criptônio

Krypton

Krypton

Criptón

криптон

קריפטון

氙
ワ
リ
プ
ト
ン

Naturally occurring isotopes: 84, 86, 82, 83, 80, 78

Density: $3.733 \times 10^{-3} \text{ g/cm}^3$ (20°C)

Melting point: -156.6°C **Boiling point:** $-152.30 \pm 0.10^\circ\text{C}$

Latent heat of fusion: 19.54 J/g

Specific heat: 0.24804 J/g/°C (25°C)

Thermal conductivity: 0.0949 mw/cm/°C (27°C)

Ionization potential (1st): 13.999 eV

Chemical valence: 0

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6$

Valence electrons: $(4s^2 4p^6)$

Crystal form: Cubic, face centered (solid)

Cross section σ : 24.5 ± 1.0 barns

La

Lanthanum

57

138.9055

IIIB

44 95592
Sc 21
88 9050
Y 39
138 9055
La 57
227 02777
Ac 89

Lantânio

Lanthane

Lanthan

Lantano

лантан

lanthan

金 蘭
ラン
タン

Naturally occurring isotopes: 139, 138**Density:** 6.145 g/cm³ (25°C)**Melting point:** 921°C **Boiling point:** 3457°C**Latent heat of fusion:** 81.4 J/g**Specific heat:** 0.195 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 5.2×10^{-6} cm/cm/°C (25°C)**Thermal conductivity:** 0.134 w/cm/°C (25°C)**Electrical resistivity:** 56×10^{-6} ohm-cm (25°C)**Ionization potential (1st):** 5.577 eV**Electron work function ϕ :** 3.5 eV**Oxidation potential:** La \rightarrow La³⁺ + 3e = 2.522 V**Chemical valence:** 3**Electrochemical equivalents:** 1.7275 g/amp-hr**Ionic radius:** 1.061 Å (La³⁺)**Valence electron potential ($-\epsilon$ V):** 40.71**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p⁶
5d¹ 6s²**Valence electrons:** 5d¹ 6s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 8.9 \pm 0.2 barns**Vapor pressure:** 1.33 \times 10⁻⁷ Pa (at melting point)

Lr

Lawrencium

103

260

Laurêncio
Lawrencium
Lawrenzium
Lawrencio
лавренций

Actinide Series

232.03807 Th 90	231.0358 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

テ
ー
レ
ン
チ
ウ
ム**Naturally occurring isotopes:** None**Oxidation potential:** $\text{Lr} \rightarrow \text{Lr}^{3+} + 3\text{e} = 2.0 \text{ V}$ **Chemical valence:** 3**Electrochemical equivalents:** 3.23 g/amp-hr**Principal quantum number:** 7**Principal electron shells:** K L M N O P Q**Electronic configuration:** $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 5f^{14} 6s^2 6p^6 6d^1 7s^2$ **Valence electrons:** $5f^{14} 6d^1 7s^2$ **Half life:** 3 minutes

Pb

Lead

82

207.2

IVA

12.011 C 6
28.0855 Si 14
72.59 Ge 32
118.69 Sn 50
207.2 Pb 82

Chumbo

Plomb

Blei

Plomo

свинец

עופרת

鉛 鉛

Naturally occurring isotopes: 208, 206, 207, 204

Density: 11.342 g/cm³ (20°C)

Melting point: 327.502°C **Boiling point:** 1740°C

Latent heat of fusion: 23.06 J/g

Specific heat: 0.128 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 28.3×10^{-6} cm/cm/°C (25°C)

Thermal conductivity: 0.353 w/cm/°C (25°C)

Electrical resistivity: 20.65×10^{-6} ohm-cm (20°C)

Ionization potential (1st): 7.416 eV

Electron work function ϕ : 4.25 eV

Oxidation potential: $\text{Pb} \rightarrow \text{Pb}^{2+} + 2e = 0.126 \text{ V}$

Chemical valence: 2, 4

Electrochemical equivalents: 3.865 g/amp-hr (Pb^{2+})

Ionic radius: 1.19 Å (Pb^{2+})

Valence electron potential ($-\epsilon\text{V}$): 24.2

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 6s^2 6p^2$

Valence electrons: $6s^2 6p^2$

Crystal form: Cubic, face centered

Cross section σ : 180 ± 10 mbarns

Vapor pressure: 4.21×10^{-7} Pa (at melting point)

Li

Lithium

3

6.941

IA

1.0079	H	1
6.941	Li	3
22.98977	Na	11
39.098	K	19
85.4678	Rb	37
132.9054	Cs	55
223.01978	Fr	87

Litio

Lithium

Lithium

Litio

ЛИТИЙ

ליתיום

 鋰
 リチウム
Naturally occurring isotopes: 7, 6**Density:** 0.534 g/cm³ (20°C)**Melting point:** 180.54°C **Boiling point:** 1342°C**Latent heat of fusion:** 430.1 J/g**Specific heat:** 3.57 J/g/°C (25°C)**Coefficient of lineal thermal expansion:** 60 × 10⁻⁶ cm/cm/°C (25°C)**Thermal conductivity:** 0.848 w/cm/°C (25°C)**Electrical resistivity:** 8.55 × 10⁻⁶ ohm-cm (0°C)**Ionization potential (1st):** 5.392 eV**Electron work function ϕ :** 2.9 eV**Oxidation potential:** Li → Li⁺ + e = 3.045 V**Chemical valence:** 1**Electrochemical equivalents:** 0.2590 g/amp-hr**Ionic radius:** 0.76 Å (Li⁺)**Valence electron potential (-eV):** 19**Principal quantum number:** 2**Principal electron shells:** K L**Electronic configuration:** 1s² 2s¹**Valence electrons:** 2s¹**Crystal form:** Cubic, body centered**Cross section σ :** 71 barns**Vapor pressure:** 1.63 × 10⁻⁸ Pa (at melting point)

Lu

Lutetium

71

174.97

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Lutécio

Lutetium

Lutetium

Lutecio

лютеций

לּוּטֵטִיּוּם

金魯

ル
テ
チ
ウ
ム**Naturally occurring isotopes:** 175, 176**Density:** 9.840 g/cm³ (25°C)**Melting point:** 1663°C **Boiling point:** 3395°C**Latent heat of fusion:** 110.1 J/g**Specific heat:** 0.154 J/g/°C (25°C)**Coefficient of lineal thermal expansion:** 12.5 × 10⁻⁶ cm/cm/°C (400°C)**Thermal conductivity:** 0.164 w/cm/°C (25°C)**Electrical resistivity:** 79.0 × 10⁻⁶ ohm-cm (25°C)**Ionization potential (1st):** 5.4259 eV**Electron work function ϕ :** 3.3 eV**Oxidation potential:** Lu → Lu³⁺ + 3e = 2.255 V**Chemical valence:** 3**Electrochemical equivalents:** 2.1760 g/amp-hr**Ionic radius:** 0.848 Å (Lu³⁺)**Valence electron potential (-eV):** 50.9**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 5d¹ 6s²**Valence electrons:** 5d¹ 6s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 75 ± 2 barns**Vapor pressure:** 2.46 × 10³ Pa (at melting point)

Mg

Magnesium

12

24.305

IIA	
9.01218	Be
4	
24.305	Mg
12	
40.08	Ca
20	
87.62	Sr
38	
137.34	Ba
56	
226.02544	Ra
88	

Magnésio
Magnésium
Magnesium
Magnesio
магний
מגנזיום
鎂
マブネシウム

Naturally occurring isotopes: 24, 26, 25

Density: 1.738 g/cm³ (20°C)

Melting point: 648.8 ± 0.5°C **Boiling point:** 1090°C

Latent heat of fusion: 368.6 J/g

Specific heat: 102 J/g°C (25°C)

Coefficient of lineal thermal expansion: 27.1 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 1.56 w/cm°C (20°C)

Electrical resistivity: 4.45 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.646 eV

Electron work function ϕ : 3.66 eV

Oxidation potential: Mg → Mg²⁺ + 2e = 2.363 V

Chemical valence: 2

Electrochemical equivalents: 0.45341 g/amp-hr

Ionic radius: 0.72 Å (Mg²⁺)

Valence electron potential (-εV): 40

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s²

Valence electrons: 3s²

Crystal form: Hexagonal, close packed

Cross section σ : 64 ± 2 mbarns

Vapor pressure: 3.61 × 10² (at melting point)

Mn

Manganese

25

54.9380

VIIIB

54.9380
Mn 25
96.906
Tc 43
186.2
Re 75
107

Manganès

Manganese

Mangan

Manganeseo

марганец

マンガン

金子
メ
ン
ガン

Naturally occurring isotope: 55

Density: 7.44 g/cm³ (20°C)

Melting point: 1244 ± 3°C **Boiling point:** 1962°C

Latent heat of fusion: 266.7 J/g

Specific heat: 0.479 J/g°C (20°C)

Coefficient of lineal thermal expansion: 22 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 78.1 mw/cm°C (25°C)

Electrical resistivity: 185 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.435 eV

Electron work function ϕ : 4.1 eV

Oxidation potential: Mn → Mn²⁺ + 2e = 1.18 V

Chemical valence: -2, -1, 0, 1, 2, 3, 4, 5, 6, 7

Electrochemical equivalents: 0.29282 g/amp-hr

Ionic radius: 0.46 Å (Mn⁷⁺)

Valence electron potential (-εV): 220

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁵ 4s²

Valence electrons: 3d⁵ 4s²

Crystal form: Cubic, face centered

Cross section σ : 13.3 ± 0.1 barns

Vapor pressure: 1.21 × 10² Pa (at melting point)

Md

Mendelevium

101

258

Mendelëvio
 Mendeleevium
 Mendeleevium
 Mendelëvio
 менделевий
 מנדלביזיום

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

金門
 멘
 デ
 レ
 ビ
 ウ
 ム

Naturally occurring isotopes: None

Ionization potential (1st): 6.58 eV

Oxidation potential: $\text{Md} \rightarrow \text{Md}^{3+} + 3\epsilon = 1.6 \text{ V}$

Chemical valence: 1, 2, 3

Electrochemical equivalents: 3.21 g/amp-hr

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6 5d^{10} 5f^{13} 6s^2 6p^6 7s^2$

Valence electrons: $5f^{13} 7s^2$

Half life: 55 days

Hg

Mercury

80

200.59

IIB

65 38 Zn 30
112 41 Cd 48
200 59 Hg 80

Mercúrio

Mercur

Quecksilber

Mercurio

ртуть

כספית

汞 水銀

Naturally occurring isotopes: 202, 200, 199, 201, 198, 204, 196

Density: 13.534 g/cm³ (25°C)

Melting point: -38.87°C **Boiling point:** 356.58°C

Latent heat of fusion: 11.46 J/g

Specific heat: 0.1395 J/g°C (liquid) (25°C)

Thermal conductivity: 0.0830 w/cm°C (25°C)

Electrical resistivity: 95.78 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 10.437 eV

Electron work function ϕ : 4.49 eV

Oxidation potential: Hg → Hg²⁺ + 2e = -0.788 V

Chemical valence: 1, 2

Electrochemical equivalents: 3.7420 g/amp-hr

Ionic radius: 1.02 Å (Hg²⁺)

Valence electron potential (-eV): 28.2

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s²

Valence electrons: 6s²

Crystal form: Rhombohedral

Cross section σ : 375 ± 5 barns

Vapor pressure: 2.00 × 10⁻⁴ Pa (at melting point)

Mo

Molybdenum

42

95.94

VIB

51 996
Cr
24
95 94
Mo
42
183 85
W
74
106

Molibdênio

Molybdène

Molybdän

Molibdeno

молибден

מוליבדן

鉬
モリブデン

Naturally occurring isotopes: 98, 96, 95, 92, 100, 97, 94

Density: 10.22 g/cm³ (20°C)

Melting point: 2617°C **Boiling point:** 4612°C

Latent heat of fusion: 288.0 J/g

Specific heat: 0.251 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 6.6 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 1.38 w/cm/°C (25°C)

Electrical resistivity: 5.2 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 7.099 eV

Electron work function ϕ : 4.6 eV

Oxidation potential: Mo → Mo³⁺ + 3e = 0.2 V

Chemical valence: 2, 3, 4, 5, 6

Electrochemical equivalents: 0.8949 g/amp-hr

Ionic radius: 0.650 Å (Mo⁴⁺)

Valence electron potential (-εV): 88.6

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d⁵ 5s¹

Valence electrons: 4d⁵ 5s¹

Crystal form: Cubic, body centered

Cross section σ : 2.65 ± 0.05 barns

Vapor pressure: 3.47 Pa (at melting point)

Nd

Neodymium

60

144.24

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Neodímio
Neodymium
Neodym
Neodimio
неодимий
ניאודימיום

釹
ネオジウム

Naturally occurring isotopes: 142, 144, 146, 143, 145, 148, 150

Density: 7.007 g/cm³ (25°C)

Melting point: 1021°C **Boiling point:** 3068°C

Latent heat of fusion: 75.47 J/g

Specific heat: 0.190 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 8.6 × 10⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.165 w/cm/°C (25°C)

Electrical resistivity: 64.0 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 5.49 eV

Electron work function ϕ : 3.2 eV

Oxidation potential: Nd → Nd³⁺ + 3e = 2.431 V

Chemical valence: 2, 3

Electrochemical equivalents: 1.7939 g/amp-hr

Ionic radius: 0.995 Å (Nd³⁺)

Valence electron potential (-eV): 43.4

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f⁴ 5s² 5p⁶ 6s²

Valence electrons: 4f⁴ 6s²

Crystal form: Hexagonal, close packed

Cross section σ : 49 ± 2 barns

Vapor pressure: 6.03 × 10⁻³ Pa (at melting point)

Ne

Neon

10

20.179

0

4 00260 He 2
20 179 Ne 10
39 948 Ar 18
83 80 Kr 36
131 30 Xe 54
222 01761 Rn 86

Neônio

Neon

Neon

Neón

неон

ניאון

氛 ネオン

Naturally occurring isotopes: 20, 22, 21

Density: $0.8999 \times 10^{-3} \text{ g/cm}^3$ (20°C)

Melting point: -248.67°C **Boiling point:** -246.048°C

Latent heat of fusion: 16.6 J/g

Specific heat: 1.0301 J/g/°C (25°C)

Thermal conductivity: 0.493 mw/cm/°C (27°C)

Ionization potential (1st): 21.564 eV

Chemical valence: 0

Principal quantum number: 2

Principal electron shells: K L

Electronic configuration: $1s^2 2s^2 2p^6$

Valence electrons: ($2s^2 2p^6$)

Crystal form: Cubic, face centered

Cross section σ : 38 ± 10 mbarns

Np

Neptunium

93

237.0482

Actinide Series

232 02807 Th 90	231 0359 Pa 91	238 029 U 92	237 0482 Np 93	244 06423 Pu 94	243 0614 Am 95	247 07038 Cm 96	247 07032 Bk 97	251 07961 Cf 98	254 08805 Es 99
257 09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Neptúnio
 Neptunium
 Neptunium
 Neptunio
 нептуний
 נפטוניום

鐳
 ネプ
 ツニ
 ウム

Naturally occurring isotopes: None

Density: 20.45 g/cm³ (25°C)

Melting point: 640 ± 1°C **Boiling point:** 3902°C

Latent heat of fusion: 46 J/g

Thermal conductivity: 63 mw/cm/°C (27°C)

Electrical resistivity: 119 × 10⁻⁶ ohm-cm (100°C)

Ionization potential (1st): 6.19 eV

Oxidation potential: Np → Np³⁺ + 3e = 1.856 V

Chemical valence: 3, 4, 5, 6, 7

Electrochemical equivalents: 1.7689 g/amp-hr

Ionic radius: 0.75 Å (Np⁵⁺)

Valence electron potential (-εV): 96

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f⁴ 6s² 6p⁶ 6d¹ 7s²

Valence electrons: 5f⁴ 6d¹ 7s²

Crystal form: Orthorhombic

Half life: 2.14 × 10⁴ years

Cross section σ: 170 ± 5 barns

Ni

Nickel

28

58.70

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
	109	

Niquel

Nickel

Nickel

Niquel

никель

ניקל

鎳
ニッケル

Naturally occurring isotopes: 58, 60, 62, 61, 64

Density: 8.902 g/cm³ (25°C)

Melting point: 1453°C **Boiling point:** 2732°C

Latent heat of fusion: 300.3 J/g

Specific heat: 0.444 J/g°C (25°C)

Coefficient of lineal thermal expansion: 13.3 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 0.909 w/cm°C (25°C)

Electrical resistivity: 6.84 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.635 eV

Electron work function ϕ : 5.15 eV

Oxidation potential: Ni → Ni²⁺ + 2e = 0.250 V

Chemical valence: 0, 1, 2, 3

Electrochemical equivalents: 1.095 g/amp-hr

Ionic radius: 0.69 Å (Ni²⁺)

Valence electron potential (-eV): 42

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d⁸ 4s²

Valence electrons: 3d⁸ 4s²

Crystal form: Cubic, face centered

Cross section σ : 4.54 ± 0.10 barns

Vapor pressure: 2.37 × 10² Pa (at melting point)

Nb

Niobium

41

92.9064

VB	
50 9415	V
23	
92 9064	Nb
41	
180 9479	Ta
73	
105	

Nióbio

Niobium

Niob

Niobio

ниобий

ניוביום

銱
ニオブ

Naturally occurring isotope: 93

Density: 8.57 g/cm³ (20°C)

Melting point: 2468 ± 10°C **Boiling point:** 4742°C

Latent heat of fusion: 288.4 J/g

Specific heat: 0.265 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 7.31 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.537 w/cm/°C (25°C)

Electrical resistivity: 14.6 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 6.88 eV

Electron work function ϕ : 4.3 eV

Oxidation potential: Nb → Nb³⁺ + 3e = 1.099 V

Chemical valence: 2, 3, 4, 5

Electrochemical equivalents: 0.69327 g/amp-hr

Ionic radius: 0.69 Å (Nb⁵⁺)

Valence electron potential (-eV): 104

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d⁴ 5s¹

Valence electrons: 4d⁴ 5s¹

Crystal form: Cubic, face centered

Cross section σ : 1.15 ± 0.05 barns

Vapor pressure: 7.55 × 10⁻² Pa (at melting point)

N

Nitrogen

7

14.0067

VA	
14.0067	N 7
30.97376	P 15
74.9216	As 33
121.75	Sb 51
208.9804	Bi 83

Nitrogénio

Azote

Stickstoff

Nitrógeno

азот

נקוד

氮 窒素

Naturally occurring isotopes: 14, 15

Density: $1.165 \times 10^{-3} \text{ g/cm}^3$ (20°C)

Melting point: -209.86°C **Boiling point:** -195.8°C

Latent heat of fusion: 51.41 J/g (N_2)

Specific heat: 1.040 J/g°C (N_2) (25°C)

Thermal conductivity: 0.2598 mw/cm°C (27°C at 1 atm)

Ionization potential (1st): 14.534 eV

Oxidation potential: $\text{N}_2 + 2\text{H}_2\text{O} \rightarrow \text{H}_2\text{N}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- = -2.65 \text{ V}$

Chemical valence: $-3, 3, 5$

Electrochemical equivalents: 0.10452 g/amp-hr

Ionic radius: 0.13 Å (N^{5+})

Valence electron potential ($-\epsilon\text{V}$): 550

Principal quantum number: 2

Principal electron shells: K L

Electronic configuration: $1s^2 2s^2 2p^3$

Valence electrons: $2s^2 2p^3$

Crystal form: Hexagonal, close packed

Cross section σ : 1.9 barns

No

Nobelium

102

259

Nobélio
Nobelium
Nobelium
Nobelio
нобелий
נובליום

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

ノ
ベ
リ
ウ
ム

Naturally occurring isotopes: None

Ionization potential (1st): 6.65 eV

Oxidation potential: $\text{No} \rightarrow \text{No}^{2+} + 2e = 2.5 \text{ V}$

Chemical valence: 2, 3

Electrochemical equivalents: 4.83 g/amp-hr

Ionic radius: 1.1 Å (est) (No^{2+})

Valence electron potential ($-\epsilon\text{V}$): (26)

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 5f^{14} 6s^2 6p^6 7s^2$

Valence electrons: $5f^{14} 7s^2$

Half life: ~59 minutes

Os

Osmium

76

190.2

VIII		
55 847 Fe 26	58 9332 Co 27	58 70 Ni 28
101 07 Ru 44	102 9055 Rh 45	106 4 Pd 46
190 2 Os 76	192 22 Ir 77	195 09 Pt 78
109		

Osmio

Osmium

Osmium

Osmio

ОСМИЙ

𐤛𐤓𐤕𐤓𐤕

 鐵
 オ
 ス
 ミ
 ウ
 ム

Naturally occurring isotopes: 192, 190, 189, 188, 187, 186, 184

Density: 22.61 g/cm³ (20°C)

Melting point: 3045 ± 30°C **Boiling point:** 5027 ± 100°C

Latent heat of fusion: 154.1 J/g

Specific heat: 0.13 J/g°C (25°C)

Coefficient of lineal thermal expansion: 6.3 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 0.876 w/cm°C (25°C)

Electrical resistivity: 9.5 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 8.7 eV

Electron work function φ: 4.83 eV

Oxidation potential: Os + 4H₂O → OsO₄ + 8H⁺ + 8e⁻ = -0.85 V

Chemical valence: 0, 1, 2, 3, 4, 5, 6, 7, 8

Electrochemical equivalents: 1.774 g/amp-hr

Ionic radius: 0.630 Å (Os⁴⁺)

Valence electron potential (-εV): 91.4

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 5d⁶ 6s²

Valence electrons: 5d⁶ 6s²

Crystal form: Hexagonal, close packed

Cross section σ: 15.3 ± 0.7 barns

Vapor pressure: 2.52 Pa (at melting point)

O

Oxygen

8

15.9994

VIA

15.9994 O 8
32.06 S 16
78.96 Se 34
127.80 Te 52
208.98243 Po 84

Oxigênio

Oxigène

Sauerstoff

Oxigeno

кислород

ᠣᠶᠢᠨ

氧 酸素

Naturally occurring isotopes: 16, 18, 17**Density:** $1.429 \times 10^{-3} \text{ g/cm}^3$ (0°C)**Melting point:** -218.4°C **Boiling point:** -182.962°C **Latent heat of fusion:** 26.17 J/g (O₂)**Specific heat:** 0.9174 J/g/°C (O₂) (25°C)**Thermal conductivity:** 0.2674 w/cm/°C (25°C at 1 atm)**Ionization potential (1st):** 13.618 eV**Oxidation potential:** $2\text{H}_2\text{O (liquid)} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^- = -1.229 \text{ V}$ **Chemical valence:** -2**Electrochemical equivalents:** 0.29847 g/amp-hr**Ionic radius:** 1.40 Å (O²⁻)**Valence electron potential (-eV):** -20.6**Principal quantum number:** 2**Principal electron shells:** K L**Electronic configuration:** $1s^2 2s^2 2p^4$ **Valence electrons:** $2s^2 2p^4$ **Crystal form:** Cubic**Cross section σ :** 0.27 mbarns

Pd

Palladium

46

106.4

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
109		

Paládio
 Palladium
 Palladium
 Paladio
 палладий
 פלדיום
 鈀
 パラジウム

Naturally occurring isotopes: 106, 108, 105, 110, 104, 102

Density: 12.023 g/cm³ (20°C)

Melting point: 1554°C **Boiling point:** 3140°C

Latent heat of fusion: 157.4 J/g

Specific heat: 0.244 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 11.67 × 10⁻⁶ cm/cm/°C (0°C)

Thermal conductivity: 0.718 w/cm/°C (25°C)

Electrical resistivity: 10.54 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 8.34 eV

Electron work function φ: 5.12 eV

Oxidation potential: Pd → Pd²⁺ + 2ε = -0.987 V

Chemical valence: 2, 3, 4

Electrochemical equivalents: 1.985 g/amp-hr

Ionic radius: 0.86 Å (Pd²⁺)

Valence electron potential (-eV): 33

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰

Valence electrons: 4d¹⁰

Crystal form: Cubic, face centered

Cross section σ: 6.0 ± 1.0 barns

Vapor pressure: 1.33 Pa (at melting point)

P

Phosphorus

15

30.97376

VA

14.0067 N 7
30.97376 P 15
74.9216 As 33
121.75 Sb 51
208.9804 Bi 83

Fósforo
Phosphore
Phosphor
Fósforo
φοσφορ

ἄσφῳδον

磷
燐

Naturally occurring isotope: 31

Density: 1.828 g/cm³ (white), 2.34 g/cm³ (red), 2.699 g/cm³ (black)
(all at 20°C)

Melting point: 44.1°C (white) **Boiling point:** 280.3°C (white)

Latent heat of fusion: 20.28 J/g (white)

Specific heat: 0.7697 J/g°C (white) (25°C)

Coefficient of linear thermal expansion: 125 × 10⁻⁶ cm/cm°C (25°C)

Thermal conductivity: 2.36 mw/cm°C (white) (25°C)

Electrical resistivity: 10¹¹ ohm-cm (white) (20°C)

Ionization potential (1st): 10.486 eV

Oxidation potential: P + 2H₂O → H₃PO₂ + H⁺ + e = 0.508 V

Chemical valence: -3, 3, 5

Electrochemical equivalents: 0.23113 g/amp-hr

Ionic radius: 0.38 Å (P⁵⁺)

Valence electron potential (-eV): 190

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p³

Valence electrons: 3s² 3p³

Crystal form: Cubic

Cross section σ: 0.19 barns

Vapor pressure: 20.8 Pa (at melting point)

Four allotropes of phosphorus have different melting points, crystal forms, colors, and electrical conductivities. The black variety has the highest electrical conductivity.

Pt

Platinum

78

195.09

VIII		
55 847 Fe 26	58 9332 Co 27	58 70 Ni 28
101 07 Ru 44	102 9055 Rh 45	106 4 Pd 46
190 2 Os 76	192 22 Ir 77	195 09 Pt 78
109		

Platina
Platine
Plátin
Platino
платина

פלטינ

鉑 白金
[プラチナ]

Naturally occurring isotopes: 195, 194, 196, 198, 192, 190

Density: 21.45 g/cm³ (20°C)

Melting point: 1773.5°C **Boiling point:** 3827 ± 100°C

Latent heat of fusion: 100.9 J/g

Specific heat: 0.133 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 9.5 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.716 w/cm/°C (25°C)

Electrical resistivity: 9.85 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 8.96 eV

Electron work function φ: 5.65 eV

Oxidation potential: Pt → Pt²⁺ + 2e = -1.2 V

Chemical valence: 2, 3, 4

Electrochemical equivalents: 1.8197 g/amp-hr

Ionic radius: 0.625 Å (Pt⁴⁺)

Valence electron potential (-eV): 92.2

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 5d⁹ 6s¹

Valence electrons: 5d⁹ 6s¹

Crystal form: Cubic, face centered

Cross section σ: 9 ± 1 barns

Vapor pressure: 3.12 × 10⁻² Pa (at melting point)

Pu Plutonium

94

244.06423

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Plutônio
Plutonium
Plutonium
Plutonio
плутоний
פלוטוניום

鉷
フリ
ト
ニ
ウ
ム

Naturally occurring isotope: 242 (trace)

Density: 19.737 g/cm³ (25°C)

Melting point: 639.5°C **Boiling point:** 3232°C

Latent heat of fusion: 11 J/g

Specific heat: 0.14 J/g°C (25°C)

Coefficient of lineal thermal expansion: 42.3 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 0.0670 w/cm°C (25°C)

Electrical resistivity: 146.45 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 6.06 eV

Oxidation potential: Pu → Pu³⁺ + 3e = 2.031 V

Chemical valence: 3, 4, 5, 6, 7

Electrochemical equivalents: 2.2765 g/amp-hr

Ionic radius: 0.887 Å (Pu⁴⁺)

Valence electron potential (-εV): 64.9

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f⁶ 6s² 6p⁶ 7s²

Valence electrons: 5f⁶ 7s²

Crystal form: Monoclinic

Half life: 8.3 × 10⁷ years

Cross section σ: 1.8 ± 0.3 barns

Po

Polonium

84

208.98243

VIA

15.9994 O 8
32.06 S 16
78.96 Se 34
127.60 Te 52
208.98243 Po 84

Polônio
Polonium
Polonium
Polonio
полоний
פולוניום
鉛 ポロニウム

Naturally occurring isotopes: None

Density: 9.20 g/cm³ (20°C)

Melting point: 254°C **Boiling point:** 962°C

Latent heat of fusion: 60.1 J/g

Specific heat: 0.13 J/g°C (25°C)

Coefficient of lineal thermal expansion: 23.5 × 10⁻⁶ cm/cm°C (20°C)

Electrical resistivity: 42 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 8.42 eV

Oxidation potential: Po → Po²⁺ + 2e = -0.65 V

Chemical valence: -2, 0, 2, 4, 6

Electrochemical equivalents: 3.8986 g/amp-hr

Ionic radius: 2.30 Å (Po²⁻)

Valence electron potential (-eV): -12.5

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p⁴

Valence electrons: 6s² 6p⁴

Crystal form: Cubic, body centered

Half life: 103 years

Vapor pressure: 1.76 × 10⁻² Pa (at melting point)

K

Potassium

19

39.098

IA

1.0079 H 1
6.941 Li 3
22.98977 Na 11
39.098 K 19
85.4678 Rb 37
132.9054 Cs 55
223.01976 Fr 87

Potássio
Potassium

Kalium

Potasio

калий

פּוֹטַאשְׁיּוּם

鉀
カリウム

Naturally occurring isotopes: 39, 41, 40

Density: 0.862 g/cm³ (20°C)

Melting point: 63.25°C **Boiling point:** 759.9°C

Latent heat of fusion: 59.33 J/g

Specific heat: 0.757 J/g°C (25°C)

Coefficient of lineal thermal expansion: 83 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 1.025 w/cm°C (25°C)

Electrical resistivity: 7.20 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 4.341 eV

Electron work function φ: 2.30 eV

Oxidation potential: K → K⁺ + e = 2.925 V

Chemical valence: 1

Electrochemical equivalents: 1.4587 g/amp-hr

Ionic radius: 1.38 Å (K⁺)

Valence electron potential (-eV): 10.4

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 4s¹

Valence electrons: 4s¹

Crystal form: Cubic, body centered

Cross section σ: 2.1 barns

Vapor pressure: 1.06 × 10⁻⁴ Pa (at melting point)

Pr

Praseodymium

59

140.9077

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Praséodímio
Praséodyne
Praseodym
Praseodimio
празеодимий

𐤀𐤒𐤁𐤍𐤁𐤄

鎳
プラセオジウム

Naturally occurring isotope: 141

Density: 6.773 g/cm³ (25°C)

Melting point: 931°C **Boiling point:** 3512°C

Latent heat of fusion: 71.3 J/g

Specific heat: 0.193 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 6.5 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.125 w/cm/°C (25°C)

Electrical resistivity: 68 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 5.42 eV

Electron work function ϕ : 2.7 eV

Oxidation potential: Pr → Pr³⁺ + 3e = 2.462 V

Chemical valence: 3, 4

Electrochemical equivalents: 1.7524 g/amp-hr

Ionic radius: 1.013 Å (Pr³⁺)

Valence electron potential (-eV): 42.64

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f³ 5s² 5p⁶ 6s²

Valence electrons: 4f³ 6s²

Crystal form: Hexagonal, close packed

Cross section σ : 3.9 ± 0.5 barns

Pm

Promethium

61

144.913

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Promécio
Prometheum
Prometheum
Promecio
прометий
טרומתיום

鉷
プロメチウム

Naturally occurring isotopes: None

Density: $7.22 \pm 0.02 \text{ g/cm}^3$ (25°C)

Melting point: $1168 \pm 6^\circ\text{C}$ **Boiling point:** 2460°C

Latent heat of fusion: 86.7 J/g

Specific heat: $0.185 \text{ J/g}^\circ\text{C}$ (25°C)

Thermal conductivity: $0.179 \text{ w/cm}^\circ\text{C}$ (25°C)

Ionization potential (1st): 5.55 eV

Oxidation potential: $\text{Pm} \rightarrow \text{Pm}^{3+} + 3\epsilon = 2.423 \text{ V}$

Chemical valence: 3

Electrochemical equivalents: 1.8022 g/amp-hr

Ionic radius: 0.979 \AA (Pm^{3+})

Valence electron potential ($-\epsilon\text{V}$): 44.1

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^5 5s^2 5p^6 6s^2$

Valence electrons: $4f^5 6s^2$

Crystal form: Hexagonal

Half life: 17.7 years

Pa Protactinium

91

231.0359

Actinide Series

232.03807 Th 90	231.0359 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	259 No 102	260 Lr 103						

Protactinio
Protactinium
Protactinium
Protactinio
протактиний

פרוטאקטיןיום

鑷
プロトアクチニウム

Naturally occurring isotope: 231 (minute quantities only)

Density: 15.37 g/cm³ (25°C)

Melting point: 1575°C

Latent heat of fusion: 65 J/g

Specific heat: 0.12 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 11.2 × 10⁻⁶ cm/cm/°C (25°C)

Ionization potential (1st): 5.89 eV

Chemical valence: 3, 4, 5

Electrochemical equivalents: 1.7240 g/amp-hr

Oxidation potential: Pa → Pa³⁺ + 3e = 1.6 V

Ionic radius: 0.78 Å (Pa⁵⁺)

Valence electron potential (-eV): 92

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f² 6s² 6p⁶ 6d¹ 7s²

Valence electrons: 5f² 6d¹ 7s²

Crystal form: Tetragonal

Half life: 3.248 × 10⁴ years

Cross section σ: 200 ± 10 barns

Ra

Radium

88

226.02544

IIA

9 01218 Be 4
24 305 Mg 12
40 08 Ca 20
87 62 Sr 38
137 34 Ba 56
226.02544 Ra 88

Rádío

Radium

Radium

Radio

радий

רדיום

鐳
ラジウム

Naturally occurring isotope: 226 (minute quantities only)

Density: 5.5 g/cm³ (extrapolated) (20°C)

Melting point: 700°C **Boiling point:** 1140°C

Latent heat of fusion: 37 J/g (est)

Specific heat: 0.120 J/g/°C (25°C)

Thermal conductivity: 0.186 w/cm/°C (20°C)

Ionization potential (1st): 5.279 eV

Oxidation potential: Ra → Ra²⁺ + 2e = 2.916 V

Chemical valence: 2

Electrochemical equivalents: 4.2165 g/amp-hr

Ionic radius: 1.43 Å (Ra²⁺)

Valence electron potential (−eV): 20.1

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p⁶ 7s²

Valence electrons: 7s²

Half life: 1622 years

Cross section σ: 20 ± 3 barns

Vapor pressure: 3.27 × 10² Pa (at melting point)

Rn

Radon

86

222.01761

O

4.00260
He 2
20.179
Ne 10
39.948
Ar 18
83.80
Kr 36
131.30
Xe 54
222.01761
Rn 86

Radônio

Radon

Radon

Radón

радон

ռոն

気 ラドン

Naturally occurring isotopes: None (radium decay product)

Density: $9.96 \times 10^{-3} \text{ g/cm}^3$ (20°C)

Melting point: -71°C **Boiling point:** -61.8°C

Latent heat of fusion: 13.1 J/g

Specific heat: 0.09362 J/g/°C (25°C)

Thermal conductivity: 0.0364 mw/cm/°C (27°C)

Ionization potential (1st): 10.748 eV

Chemical valence: 0

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 6s^2 6p^6$

Valence electrons: (6s² 6p⁶)

Crystal form: Cubic, face centered

Half life: 3.824 days

Cross section σ : 0.72 ± 0.07 barns

Re

Rhenium

75

186.2

VIIB

54 9380
Mn 25
96 906
Tc 43
186.2
Re 75
107

Rênio

Rhenium

Rhenium

Renio

рений

רניום

 銱
レニウム
Naturally occurring isotopes: 187, 185**Density:** 21.04 g/cm³ (20°C)**Melting point:** 3180°C **Boiling point:** 5627°C (est)**Latent heat of fusion:** 177.6 J/g**Specific heat:** 0.137 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 6.7 × 10⁻⁶ cm/cm/°C (25°C)**Thermal conductivity:** 0.480 w/cm/°C (25°C)**Electrical resistivity:** 19.3 × 10⁻⁶ ohm-cm (20°C)**Ionization potential (1st):** 7.88 eV**Electron work function ϕ :** 4.96 eV**Oxidation potential:** $\text{Re} + 2\text{H}_2\text{O} \rightarrow \text{ReO}_2 + 4\text{H}^+ + 4\text{e}^- = -0.2513 \text{ V}$ **Chemical valence:** 0, 1, 2, 3, 4, 5, 6, 7**Electrochemical equivalents:** 0.9924 g/amp-hr**Ionic radius:** 0.56 Å (Re⁷⁺)**Valence electron potential ($-\epsilon\text{V}$):** 180**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 5d⁵ 6s²**Valence electrons:** 5d⁵ 6s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 85 ± 5 barns**Vapor pressure:** 3.24 Pa (at melting point)

Rh

Rhodium

45

102.9055

VIII		
55.847 Fe 26	58.9332 Co 27	58.70 Ni 28
101.07 Ru 44	102.9055 Rh 45	106.4 Pd 46
190.2 Os 76	192.22 Ir 77	195.09 Pt 78
	109	

Ródio

Rhodium

Rhodium

Rodio

родий

רודיום

銠
ロジウム

Naturally occurring isotope: 103

Density: 12.41 g/cm³ (20°C)

Melting point: 1966 ± 3°C **Boiling point:** 3727 ± 100°C

Latent heat of fusion: 211.6 J/g

Specific heat: 0.24 J/g°C (25°C)

Coefficient of lineal thermal expansion: 8.3 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 1.50 w/cm°C (25°C)

Electrical resistivity: 4.51 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.46 eV

Electron work function ϕ : 4.98 eV

Oxidation potential: Rh → Rh³⁺ + 3e = -0.80 V

Chemical valence: 2, 3, 4, 5, 6

Electrochemical equivalents: 1.2798 g/amp-hr

Ionic radius: 0.68 Å (Rh³⁺)

Valence electron potential (-εV): 64

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d⁸ 5s¹

Valence electrons: 4d⁸ 5s¹

Crystal form: Cubic, face centered

Cross section σ : 150 ± 5 barns

Vapor pressure: 6.33 × 10⁻¹ Pa (at melting point)

Rb

Rubidium

37

85.4678

IA
1 0079 H 1
6.941 Li 3
22.98977 Na 11
39.098 K 19
85.4678 Rb 37
132.9054 Cs 55
223.01976 Fr 87

Rubidio
 Rubidium
 Rubidium
 Rubidio
 рубидий
 רובידיום

銣 ルビジウム

Naturally occurring isotopes: 85, 87

Density: 1.532 g/cm³ (20°C)

Melting point: 38.89°C **Boiling point:** 686°C

Latent heat of fusion: 27.43 J/g

Specific heat: 0.3634 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 90 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.582 w/cm/°C (25°C)

Electrical resistivity: 12.84 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 4.177 eV

Electron work function ϕ : 2.16 eV

Oxidation potential: Rb → Rb⁺ + ϵ = 2.925 V

Chemical valence: 1

Electrochemical equivalents: 3.1888 g/amp-hr

Ionic radius: 1.52 Å (Rb⁺)

Valence electron potential (- ϵ V): 9.47

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 5s¹

Valence electrons: 5s¹

Crystal form: Cubic, body centered

Cross section σ : 0.5 ± 0.1 barns

Vapor pressure: 1.56 × 10⁻⁴ Pa (at melting point)

Ru

Ruthenium

44

101.07

VIII		
55 847 Fe 26	58 9332 Co 27	58 70 Ni 28
101 07 Ru 44	102 9055 Rh 45	106 4 Pd 46
190 2 Os 76	192 22 Ir 77	195 08 Pt 78
	109	

Rutênio
Ruthénium
Ruthenium
Rutenio
рутений
רותניום
釘 ルテニウム

Naturally occurring isotopes: 102, 104, 101, 99, 100, 96, 98

Density: 12.45 g/cm³ (20°C)

Melting point: 2310°C **Boiling point:** 3900°C

Latent heat of fusion: 252.7 J/g

Specific heat: 0.238 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 9.91×10^{-6} cm/cm/°C (50°C)

Thermal conductivity: 1.17 w/cm/°C (25°C)

Electrical resistivity: 6.80×10^{-6} ohm-cm (0°C)

Ionization potential (1st): 7.37 eV

Electron work function ϕ : 4.71 eV

Oxidation potential: $\text{Ru} + 5\text{Cl}^- \rightarrow \text{RuCl}_3^{2-} + 3\text{e} = -0.601 \text{ V}$

Chemical valence: 1, 2, 3, 4, 5, 6, 7, 8

Electrochemical equivalents: 1.2570 g/amp-hr

Ionic radius: 0.68 Å (Ru⁴⁺)

Valence electron potential (-eV): 64

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^7 5s^1$

Valence electrons: $4d^7 5s^1$

Crystal form: Hexagonal, close packed

Cross section σ : 3.0 ± 0.8 barns

Vapor pressure: 1.40 Pa (at melting point)

Sm

Samarium

62

150.4

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Samário
Samarium
Samarium
Samario
самарий
סמריום

釷
サ
マ
リ
ウ
ム

Naturally occurring isotopes: 152, 154, 147, 149, 148, 150, 144

Density: 7.520 g/cm³ (25°C)

Melting point: 1077°C **Boiling point:** 1791°C

Latent heat of fusion: 73.8 J/g

Specific heat: 0.196 J/g°C (25°C)

Thermal conductivity: 0.133 w/cm/°C (25°C)

Electrical resistivity: 88 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 5.63 eV

Electron work function ϕ : 2.7 eV

Oxidation potential: Sm → Sm³⁺ + 3e = 2.414 V

Chemical valence: 2, 3

Electrochemical equivalents: 1.870 g/amp-hr

Ionic radius: 0.964 Å (Sm³⁺)

Valence electron potential (-eV): 44.8

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f⁶ 5s²
5p⁶ 6s²

Valence electrons: 4f⁶ 6s²

Crystal form: Rhombohedral

Cross section σ : 5820 ± 100 barns

Vapor pressure: 5.63 × 10² Pa (at melting point)

Sc

Scandium

21

44.95592

IIIB

44 95592
Sc 21
86 9059
Y 39
138 9055
La 57
227 02777
Ac 89

Escândio

Scandium

Scandium

Escandio

скандий

סקנדיום

鈮
スカ
ン
ジ
ウム

Naturally occurring isotope: 45**Density:** 2.989 g/cm³ (25°C)**Melting point:** 1541°C **Boiling point:** 2831°C**Latent heat of fusion:** 358.6 J/g**Specific heat:** 0.568 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 12 × 10⁻⁶ cm/cm°C (25°C)**Thermal conductivity:** 0.158 w/cm°C (25°C)**Electrical resistivity:** 61.0 × 10⁻⁶ ohm-cm (20°C)**Ionization potential (1st):** 6.54 eV**Electron work function ϕ :** 3.5 eV**Oxidation potential:** Sc → Sc³⁺ + 3e = 2.077 V**Chemical valence:** 3**Electrochemical equivalents:** 0.55914 g/amp-hr**Ionic radius:** 0.745 Å (Sc³⁺)**Valence electron potential (-eV):** 58.0**Principal quantum number:** 4**Principal electron shells:** K L M N**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹ 4s²**Valence electrons:** 3d¹ 4s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 25 ± 2 barns**Vapor pressure:** 2.21 × 10 Pa (at melting point)

Se

Selenium

34

78.96

VIA

15 9994 O 8
32.06 S 16
78.96 Se 34
127.60 Te 52
208.98243 Po 84

Selênio
 Sélénium
 Selen
 Selenio
 селен
 סלן
 硒 セレン

Naturally occurring isotopes: 80, 78, 82, 76, 77, 74

Density: 4.792 g/cm³ (gray) (20°C)

Melting point: 217°C (gray) **Boiling point:** 684.9 ± 1.0°C

Latent heat of fusion: 68.93 J/g

Specific heat: 0.1606 J/g°C (Se₂) (25°C)

Coefficient of lineal thermal expansion: 36.8 cm/cm°C (20°C)

Thermal conductivity: 0.0452 w/cm°C (along C-axis at 25°C)

Electrical resistivity: 1 ohm-cm (20°C)

Ionization potential (1st): 9.752 eV

Electron work function ϕ : 5.9 eV

Oxidation potential: $\text{Se} + 3\text{H}_2\text{O} \rightarrow \text{H}_2\text{SeO}_3 + 4\text{H}^+ + 4\text{e}^- = -0.740 \text{ V}$

Chemical valence: -2, 4, 6

Electrochemical equivalents: 0.73650 g/amp-hr

Ionic radius: 0.50 Å (Se⁴⁺)

Valence electron potential (-eV): 120

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁴

Valence electrons: 4s² 4p⁴

Crystal forms: Hexagonal, monoclinic, amorphous

Cross section σ : 12.2 ± 0.6 barns

Vapor pressure: 6.95 × 10⁻¹ Pa (at melting point)

Si

Silicon

14

28.0855

IVA

12 011
C 6
28 0855
Si 14
72 59
Ge 32
118 69
Sn 50
207 2
Pb 82

Silicio
Silicium
Silizium
Silicio
кремний
צורן

硅 珪素

Naturally occurring isotopes: 28, 29, 30

Density: 2.329 g/cm³ (25°C)

Melting point: 1410°C **Boiling point:** 2355°C

Latent heat of fusion: 1.655 J/g

Specific heat: 0.712 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 4.2×10^{-6} cm/cm/°C (25°C)

Thermal conductivity: 1.49 w/cm/°C (25°C)

Electrical resistivity: 3.5 ohm-cm (20°C)

Ionization potential (1st): 8.151 eV

Electron work function ϕ : 4.52 eV

Oxidation potential: $\text{Si} + 2\text{H}_2\text{O} \rightarrow \text{SiO}_2 + 4\text{H}^+ + 4\text{e}^- = 0.857 \text{ V}$

Chemical valence: -4, -1, 1, 4

Electrochemical equivalents: 0.26197 g/amp-hr

Ionic radius: 0.400 Å (Si⁴⁺)

Valence electron potential (-eV): 144

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p²

Valence electrons: 3s² 3p²

Crystal form: Cubic, diamond

Cross section σ : 160 ± 20 mbarns

Vapor pressure: 4.77 Pa (at melting point)

Ag

Silver

47

107.868

IB

63 546
Cu 29
107 868
Ag 47
196 965
Au 79

Prata

Argent

Silber

Plata

серебро

᠑᠓

銀 銀

Naturally occurring isotopes: 107, 109

Density: 10.50 g/cm³ (20°C)

Melting point: 961.93°C **Boiling point:** 2212°C

Latent heat of fusion: 104.8 J/g

Specific heat: 0.2350 J/g°C (25°C)

Coefficient of lineal thermal expansion: 18.62 × 10⁻⁶ cm/cm°C (17°C)

Thermal conductivity: 4.29 w/cm°C (25°C)

Electrical resistivity: 1.586 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.576 eV

Electron work function ϕ : 4.26 eV

Oxidation potentials: Ag → Ag⁺ + ϵ = -0.7991 V

Ag⁺ → Ag²⁺ + ϵ = -1.980 V

Chemical valence: 1, 2, 3

Electrochemical equivalents: 4.0246 g/amp-hr

Ionic radius: 1.26 Å (Ag⁺)

Valence electron potential (- ϵ V): 11.4

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s¹

Valence electrons: (4d¹⁰) 5s¹

Crystal form: Cubic, face centered

Cross section σ : 63.8 ± 0.6 barns

Vapor pressure: 3.42 × 10⁻¹ Pa (at melting point)

Na

Sodium

11

22.98977

IA

1.0079 H 1
6.941 Li 3
22.98977 Na 11
39.098 K 19
85.4678 Rb 37
132.9054 Cs 55
223.01976 Fr 87

Sódio
Sodium
Natrium
Sodio
натрий
נתרן

鈉 ナトリウム

Naturally occurring isotopes: 23

Density: 0.9712 g/cm³ (20°C)

Melting point: 97.81 ± 0.03°C **Boiling point:** 882.9°C

Latent heat of fusion: 113 J/g

Specific heat: 1.23 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 72 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 1.42 w/cm/°C (25°C)

Electrical resistivity: 4.33 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 5.139 eV

Electron work function ϕ : 2.75 eV

Oxidation potential: Na → Na⁺ + e = 2.714 V

Chemical valence: 1

Electrochemical equivalents: 0.85775 g/amp-hr

Ionic radius: 1.02 Å (Na⁺)

Valence electron potential (-eV): 14.1

Principal quantum number: 3

Principal electron shells: K L M

Electronic configuration: 1s² 2s² 2p⁶ 3s¹

Valence electrons: 3s¹

Crystal form: Cubic, body centered

Cross section σ : 534 ± 5 mbarns

Vapor pressure: 1.43 × 10⁻⁵ Pa (at melting point)

Sr

Strontium

38

87.62

IIA

9 01218
Be 4
24 305
Mg 12
40 08
Ca 20
87 62
Sr 38
137 34
Ba 56
226 02544
Ra 88

Estrôncio

Strontium

Strontium

Estroncio

стронций

סטרונציום

銩
ストロンチウム

Naturally occurring isotopes: 88, 86, 87, 84

Density: 2.54 g/cm³ (20°C)

Melting point: 769°C **Boiling point:** 1384°C

Latent heat of fusion: 105.1 J/g

Specific heat: 0.30 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 21 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.354 w/cm/°C (25°C)

Electrical resistivity: 23 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 5.695 eV

Electron work function ϕ : 2.59 eV

Oxidation potential: Sr → Sr²⁺ + 2e = 2.888 V

Chemical valence: 2

Electrochemical equivalents: 1.635 g/amp-hr

Ionic radius: 1.12 Å (Sr²⁺)

Valence electron potential (-εV): 25.7

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 5s²

Valence electrons: 5s²

Crystal form: Cubic, face centered

Cross section σ : 1.21 ± 0.06 barns

Vapor pressure: 2.46 × 10² Pa (at melting point)

S

Sulfur

16

32.06

VIA

15 9994 O 8
32 06 S 16
78 96 Se 34
127 60 Te 52
208 98243 Po 84

Enxófre

Soufre

Schwefel

Azufre

cepa

גופרית

硫 硫黄

Naturally occurring isotopes: 32, 34, 33, 36**Density:** 2.07 g/cm³ (rhombic form at 25°C)**Melting point:** 112.8°C **Boiling point:** 444.674°C**Latent heat of fusion:** 44.01 J/g**Specific heat:** 0.706 J/g°C (rhombic) (25°C)**Coefficient of lineal thermal expansion:** 64.13 × 10⁻⁶ cm/cm/°C (20°C)**Thermal conductivity:** 2.70 mw/cm/°C (25°C)**Electrical resistivity:** 2 × 10¹⁷ ohm-cm (20°C)**Ionization potential (1st):** 10.360 eV**Oxidation potentials:** S + 3H₂O → H₂SO₃ + 4H⁺ + 4e⁻ = -0.45 VS²⁻ → S + 2e⁻ = 0.447 V**Chemical valence:** -2, 4, 6**Electrochemical equivalents:** 0.2990 g/amp-hr**Ionic radius:** 0.37 Å (S⁴⁺)**Valence electron potential (-eV):** 160**Principal quantum number:** 3**Principal electron shells:** K L M**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁴**Valence electrons:** 3s² 3p⁴**Crystal form:** Orthorhombic**Cross section σ:** 0.51 barns**Vapor pressure:** 2.65 × 10⁻²⁰ Pa (at melting point)

Ta

Tantalum

73

180.9479

VB

50 9415
V
23
52 9064
Nb
41
180 9479
Ta
73
105

Tantálio

Tantale

Tantal

Tántalo

тантал

טנטל

鉭
タニブム

Naturally occurring isotopes: 181, 180

Density: 16.60 g/cm³ (20°C)

Melting point: 2996°C **Boiling point:** 5425 ± 100°C

Latent heat of fusion: 174 J/g

Specific heat: 0.140 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 6.5 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.575 w/cm/°C (25°C)

Electrical resistivity: 12.45 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 7.89 eV

Electron work function ϕ : 4.25 eV

Oxidation potential: 2Ta + 5H₂O → Ta₂O₅ + 10H⁺ + 10e = 0.812 V

Chemical valence: 3, 4, 5

Electrochemical equivalents: 1.3502 g/amp-hr

Ionic radius: 0.64 Å (Ta⁵⁺)

Valence electron potential (-εV): 110

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d³ 6s²

Valence electrons: 5d³ 6s²

Crystal form: Cubic, body centered

Cross section σ : 22 ± 1 barns

Vapor pressure: 7.76 × 10⁻¹ Pa (at melting point)

Tc

Technetium

43

96.906

VII B

54 3380
Mn 25
96 906
Tc 43
186 2
Re 75
107

Tecnécio

Technetium

Technetium

Tecnecio

технеций

טכנציום

錳
テ
ク
ネ
チ
ウム

Naturally occurring isotopes: None

Density: 11.496 g/cm³ (25°C)

Melting point: 2172°C **Boiling point:** 4877°C

Latent heat of fusion: 235 ± 5 J/g

Specific heat: 0.24 J/g/°C (25°C)

Thermal conductivity: 0.506 w/cm/°C (25°C)

Ionization potential (1st): 7.28 eV

Oxidation potential: Tc → Tc²⁺ + 2e = -0.4 V

Chemical valence: 0, 1, 2, 3, 4, 5, 6, 7

Electrochemical equivalents: 0.51651 g/amp-hr

Ionic radius: 0.56 Å (Tc⁷⁺)

Valence electron potential (-εV): 180

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d⁶ 5s¹

Valence electrons: 4d⁶ 5s¹

Crystal form: Hexagonal, close packed

Half life: 2.6 × 10⁶ years

Vapor pressure: 2.29 × 10⁻² Pa (at melting point)

Te

Tellurium

52

127.60

VIA

15 9994 O 8
32 06 S 16
78 96 Se 34
127 60 Te 52
208 98243 Po 84

Telúrio

Tellure

Tellur

Telurio

טלוריום

טלור

碲
テルル**Naturally occurring isotopes:** 130, 128, 126, 125, 124, 122, 123**Density:** 6.24 g/cm³ (20°C)**Melting point:** 449.5 ± 0.3°C **Boiling point:** 989.8 ± 3.8°C**Latent heat of fusion:** 137.2 J/g**Specific heat:** 0.202 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 16.75 × 10⁻⁶ cm/cm°C (20°C)**Thermal conductivity:** 0.0338 w/cm°C (along C-axis at 25°C)**Electrical resistivity:** 4.36 ohm-cm (25°C)**Ionization potential (1st):** 0.009 eV**Electron work function ϕ :** 4.95 eV**Oxidation potential:** $\text{Te} + 2\text{H}_2\text{O} \rightarrow \text{TeO}_2 + 4\text{H}^+ + 4\text{e}^- = -0.529 \text{ V}$ **Chemical valence:** -2, 2, 4, 6**Electrochemical equivalents:** 1.1902 g/amp-hr**Ionic radius:** 0.97 Å (Te⁴⁺)**Valence electron potential (- ϵ V):** 59**Principal quantum number:** 5**Principal electron shells:** K L M N O**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p⁴**Valence electrons:** 5s² 5p⁴**Crystal form:** Hexagonal**Cross section σ :** 4.7 ± 0.1 barns**Vapor pressure:** 2.31 × 10 Pa (at melting point)

Tb

Terbium

65

158.9254

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Térbio
Terbium
Terbium
Terbio
тербий

טרביום

銲
テルビウム

Naturally occurring isotope: 159

Density: 8.229 g/cm³ (25°C)

Melting point: 1356°C **Boiling point:** 3123°C

Latent heat of fusion: 102.7 J/g

Specific heat: 0.182 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 11.8 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.111 w/cm/°C (25°C)

Electrical resistivity: 116 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 5.85 eV

Electron work function ϕ : 3.0 eV

Oxidation potential: Tb → Tb³⁺ + 3e = 2.391 V

Chemical valence: 3, 4

Electrochemical equivalents: 1.9765 g/amp-hr

Ionic radius: 0.923 Å (Tb³⁺)

Valence electron potential (-εV): 46.8

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f⁹ 5s² 5p⁶ 6s²

Valence electrons: 4f⁹ 6s²

Crystal form: Hexagonal, close packed

Cross section σ : 30 ± 10 barns

Tl

Thallium

81

204.37

IIIA	
10.81 B 5	
26.98154 Al 13	
69.72 Ga 31	
114.82 In 49	
204.37 Tl 81	

Tálio
Thallium
Thallium
Talio
таллий
תליום

鈊
タリウム

Naturally occurring isotopes: 205, 203

Density: 11.85 g/cm³ (20°C)

Melting point: 303.5°C **Boiling point:** 1457 ± 10°C

Latent heat of fusion: 20.90 J/g

Specific heat: 0.129 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 28 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.461 w/cm/°C (25°C)

Electrical resistivity: 18.0 × 10⁻⁶ ohm-cm (0°C)

Ionization potential (1st): 6.108 eV

Electron work function ϕ : 3.84 eV

Oxidation potentials: Tl → Tl⁺ + ϵ = 0.3363 V

Tl⁺ → Tl³⁺ + 2 ϵ = -1.25 V

Chemical valence: 7, 3

Electrochemical equivalents: 7.6250 g/amp-hr

Ionic radius: 1.50 Å (Tl⁺)

Valence electron potential (- ϵ V): 9.60

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p¹

Valence electrons: 6s² 6p¹

Crystal form: Hexagonal, close packed

Cross section σ : 3.4 ± 0.5 barns

Vapor pressure: 5.33 × 10⁻⁶ Pa (at melting point)

Th

Thorium

90

232.03807

Actinide Series

232.03807	231.0358	238.029	237.0482	244.06423	243.0614	247.07038	247.07032	251.07961	254.08805
Th 90	Pa 91	U 92	Np 93	Pu 94	Am 95	Cm 96	Bk 97	Cf 98	Es 99
257.09515	258	259	260						
Fm 100	Md 101	No 102	Lr 103						

Tório
 Thorium
 Thorium
 Torio
 торий
 תוריום

釷 トリウム

Naturally occurring isotope: 232

Density: 11.724 g/cm³ (25°C)

Melting point: 1750°C **Boiling point:** 4787°C

Latent heat of fusion: 82.93 J/g

Specific heat: 0.118 J/g°C (25°C)

Coefficient of lineal thermal expansion: 12.5 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.540 w/cm/°C (25°C)

Electrical resistivity: 13.1 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.08 eV

Electron work function ϕ : 3.41 eV

Oxidation potential: Th → Th⁴⁺ + 4e = 1.899 V

Chemical valence: 3, 4

Electrochemical equivalents: 2.1643 g/amp-hr

Ionic radius: 0.972 Å (Th⁴⁺)

Valence electron potential (-eV): 59.3

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 6s² 6p⁶ 6d² 7s²

Valence electrons: 6d² 7s²

Crystal form: Cubic, face centered

Half life: 1.40 × 10¹⁰ years

Cross section σ : 74 ± 0.1 barns

Tm

Thulium

69

168.9342

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Túlio
Thulium
Thulium
Tulio
тулий
תוליום

鈹
ツリウム

Naturally occurring isotope: 169

Density: 9.321 g/cm³ (25°C)

Melting point: 1545 ± 15°C **Boiling point:** 1727°C

Latent heat of fusion: 109.0 J/g

Specific heat: 0.160 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 11.6 × 10⁻⁶ cm/cm/°C (400°C)

Thermal conductivity: 0.169 w/cm/°C (25°C)

Electrical resistivity: 79 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.1844 eV

Oxidation potential: Tm → Tm³⁺ + 3e = 2.278 V

Chemical valence: 2, 3

Electrochemical equivalents: 2.1010 g/amp-hr

Ionic radius: 0.869 Å (Tm³⁺)

Valence electron potential (-εV): 49.7

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹³ 5s² 5p⁶ 6s²

Valence electrons: 4f¹³ 6s²

Crystal form: Hexagonal, close packed

Cross section σ: 115 ± 15 barns

Vapor pressure: 4.90 × 10⁻³ Pa (at melting point)

Sn

Tin

50

118.69

IVA

12 011
C
6
28 0855
Si
14
72 59
Ge
32
118 69
Sn
50
207 2
Pb
82

Estanho

Etain

Zinn

Estaño

оолово

ברזל

錫 すす

Naturally occurring isotopes: 120, 118, 116, 119, 117, 124, 122, 112, 114, 115

Density: 7.298 g/cm³ (25°C)

Melting point: 231.9681°C **Boiling point:** 2270°C

Latent heat of fusion: 60.67 J/g

Specific heat: 0.227 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 23 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 0.668 w/cm/°C (25°C)

Electrical resistivity: 11.5 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 7.334 eV

Electron work function ϕ : 4.42 eV

Oxidation potentials: Sn → Sn²⁺ + 2e = 0.136 V

Sn²⁺ → Sn⁴⁺ + 2e = -0.15 V

Chemical valence: -4, -1, 2, 4

Electrochemical equivalents: 1.1071 g/amp-hr

Ionic radius: 0.690 Å (Sn⁴⁺)

Valence electron potential (-eV): 83.5

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 5s² 5p²

Valence electrons: 5s² 5p²

Crystal form: Tetragonal

Cross section σ : 0.63 ± 0.1 barns

Vapor pressure: 5.78 × 10⁻²¹ Pa (at melting point)

Ti

Titanium

22

47.90

IVB

47.90 Ti 22
91.22 Zr 40
178.49 Hf 72
104

Titânio

Titane

Titan

Titanio

ТИТАН

טִיטַנְיָוּם

 鈦
チタン
Naturally occurring isotopes: 48, 46, 47, 49, 50**Density:** 4.507 g/cm³ (20°C)**Melting point:** 1660 ± 10°C **Boiling point:** 3287°C**Latent heat of fusion:** 323.4 J/g**Specific heat:** 0.522 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 8.41 × 10⁻⁶ cm/cm°C (20°C)**Thermal conductivity:** 0.219 w/cm°C (25°C)**Electrical resistivity:** 42 × 10⁻⁶ ohm-cm (20°C)**Ionization potential (1st):** 6.82 eV**Electron work function ϕ :** 4.33 eV**Oxidation potential:** Ti → Ti²⁺ + 2e = 1.628 V**Chemical valence:** 1, 2, 3, 4**Electrochemical equivalents:** 0.4468 g/amp-hr**Ionic radius:** 0.605 Å (Ti⁴⁺)**Valence electron potential (-eV):** 95.2**Principal quantum number:** 4**Principal electron shells:** K L M N**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d² 4s²**Valence electrons:** 3d² 4s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 6.1 ± 0.2 barns**Vapor pressure:** 4.90 × 10⁻¹ Pa (at melting point)

W

Tungsten

74

183.85

VIB

51 996
Cr 24
95 94
Mo 42
183 85
W 74
106

Tungstênio

Tungstène

Wolframz

Tungsteno

вольфрам

וולטרם

 錫
 ツ
 ン
 ク
 ス
 テ
 ン
Naturally occurring isotopes: 184, 186, 182, 183, 180**Density:** 19.35 g/cm³ (20°C)**Melting point:** 3410 ± 20°C **Boiling point:** 5660°C**Latent heat of fusion:** 191.7 J/g**Specific heat:** 0.125 J/g/°C (25°C)**Coefficient of linear thermal expansion:** 4.6 × 10⁻⁶ cm/cm/°C (20°C)**Thermal conductivity:** 1.73 w/cm/°C (25°C)**Electrical resistivity:** 5.65 × 10⁻⁶ ohm-cm (27°C)**Ionization potential (1st):** 7.98 eV**Electron work function ϕ :** 4.55 eV**Oxidation potential:** $W + 3H_2O \rightarrow WO_3 + 6H^+ + 6e = 0.09 V$ **Chemical valence:** 2, 3, 4, 5, 6**Electrochemical equivalents:** 1.1432 g/amp-hr**Ionic radius:** 0.62 Å (W⁶⁺)**Valence electron potential (-eV):** 140**Principal quantum number:** 6**Principal electron shells:** K L M N O P**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 5d⁴ 6s²**Valence electrons:** 5d⁴ 6s²**Crystal form:** Alpha—cubic, body centered; beta—cubic, face centered**Cross section σ :** 18.5 ± 0.5 barns**Vapor pressure:** 4.27 Pa (at melting point)

U

Uranium

92

238.029

Actinide Series

232.03807 Th 90	231.0358 Pa 91	238.029 U 92	237.0482 Np 93	244.06423 Pu 94	243.0614 Am 95	247.07038 Cm 96	247.07032 Bk 97	251.07961 Cf 98	254.08805 Es 99
257.09515 Fm 100	258 Md 101	258 No 102	260 Lr 103						

Urânio
Uranium
Uran
Uranio
уран

אורניום

鈾
ウ
ラ
ニ
ウ
ム

Naturally occurring isotopes: 238, 235, 234

Density: 19.04 g/cm³ (25°C)

Melting point: 1132.3 ± 0.8°C Boiling point: 3818°C

Latent heat of fusion: 65.08 J/g

Specific heat: 0.1162 J/g°C (25°C)

Coefficient of lineal thermal expansion: 13.4 × 10⁻⁶ cm/cm°C (25°C)

Thermal conductivity: 0.275 w/cm°C (25°C)

Electrical resistivity: 27 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.05 eV

Electron work function φ: 3.63 eV

Oxidation potential: U → U³⁺ + 3e = 1.789 V

Chemical valence: 3, 4, 5, 6

Electrochemical equivalents: 1.4801 g/amp-hr

Ionic radius: 0.52 Å (U⁶⁺)

Valence electron potential (-eV): 170

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electron configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶
5d¹⁰ 5f³ 6s² 6p⁶ 6d¹ 7s²Valence electrons: 5f³ 6d¹ 7s²

Crystal form: Orthorhombic

Half life: 4.51 × 10⁹ years

Cross section σ: 7.595 ± 0.070 barns

Vapor pressure σ: 1.19 × 10⁻⁶ Pa (at melting point)

V

Vanadium

23

50.9415

VB	
50 9415	V 23
92 9064	Nb 41
180 9479	Ta 73
105	

Vanádio
Vanadium
Vanadium
Vanadio
ванадий
𐤅𐤆𐤃𐤅
𐤅𐤆𐤃𐤅
𐤅𐤆𐤃𐤅

Naturally occurring isotopes: 51, 50

Density: 6.11 g/cm³ (18.7°C)

Melting point: 1890 ± 10°C **Boiling point:** 3380°C

Latent heat of fusion: 345.2 J/g

Specific heat: 0.489 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 6.15 × 10⁻⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.307 w/cm/°C (25°C)

Electrical resistivity: 24.8 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 6.74 eV

Electron work function ϕ : 4.3 eV

Oxidation potential: V → V²⁺ + 2e = 1.186 V

Chemical valence: 2, 3, 4, 5

Electrochemical equivalents: 0.38013 g/amp-hr

Ionic radius: 0.59 Å (V⁵⁺)

Valence electron potential (-eV): 120

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d³ 4s²

Valence electrons: 3d³ 4s²

Crystal form: Cubic, body centered

Cross section σ : 5.06 ± 0.06 barns

Vapor pressure: 3.06 Pa (at melting point)

Xe

Xenon

54

131.30

O	
4 00260	He 2
20 179	Ne 10
39 948	Ar 18
83 80	Kr 36
131 30	Xe 54
222.01761	Rn 86

Хелônio

Хénon

Хелон

Хелòн

ксéнон

ქსéონი

氙 キシノン

Naturally occurring isotopes: 132, 129, 131, 134, 136, 130, 128, 124, 126

Density: $5.895 \times 10^{-3} \text{ g/cm}^3$ (20°C)

Melting point: -111.9°C **Boiling point:** $-107.1 \pm 3^\circ\text{C}$

Latent heat of fusion: 17.5 J/g

Specific heat: 0.15831 J/g°C (25°C)

Thermal conductivity: 0.514 mW/cm°C (0°C at 1 atm)

Ionization potential (1st): 12.130 eV

Chemical valence: 0

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 5s^2 5p^6$

Valence electrons: ($5s^2 5p^6$)

Crystal form: Cubic, face centered

Cross section σ : 24.5 ± 1.0 barns

Yb

Ytterbium

70

173.04

Lanthanide Series

140.12 Ce 58	140.9077 Pr 59	144.24 Nd 60	144.913 Pm 61	150.4 Sm 62	151.96 Eu 63	157.25 Gd 64	158.9254 Tb 65	162.50 Dy 66	164.9304 Ho 67
167.26 Er 68	168.9342 Tm 69	173.04 Yb 70	174.97 Lu 71						

Itérbio
Ytterbium
Ytterbium
Itterbio
иттербий
איטרביום

鏡
イ
ッ
テ
ル
ビ
ウ
ム

Naturally occurring isotopes: 174, 172, 173, 171, 176, 170, 168

Density: 6.965 g/cm³ (25°C)

Melting point: 819°C **Boiling point:** 1194°C

Latent heat of fusion: 53.23 J/g

Specific heat: 0.155 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 29.9 × 10⁶ cm/cm/°C (25°C)

Thermal conductivity: 0.349 w/cm/°C (25°C)

Electrical resistivity: 28 × 10⁻⁶ ohm-cm (25°C)

Ionization potential (1st): 6.2539 eV

Oxidation potential: Yb → Yb³⁺ + 3e = 2.267 V

Chemical valence: 2, 3

Electrochemical equivalents: 2.1520 g/amp-hr

Ionic radius: 0.858 Å (Yb³⁺)

Valence electron potential (-eV): 50.3

Principal quantum number: 6

Principal electron shells: K L M N O P

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹⁰ 4f¹⁴ 5s² 5p⁶ 6s²

Valence electrons: 4f¹⁴ 6s²

Crystal form: Cubic, face centered

Cross section σ: 37 ± 3 barns

Vapor pressure: 3.95 × 10² Pa (at melting point)

Y

Yttrium

39

88.9059

IIIB

44 95592
Sc 21
88 9059
Y 39
138 9055
La 57
227 02777
Ac 89

Itrio

Yttrium

Yttrium

Itrio

ИТТРИЙ

איטריום

 釷
 イ
 ッ
 ト
 リ
 ム
Naturally occurring isotope: 89**Density:** 4.469 g/cm³ (25°C)**Melting point:** 1522°C **Boiling point:** 3338°C**Latent heat of fusion:** 193.1 J/g**Specific heat:** 0.298 J/g°C (25°C)**Coefficient of lineal thermal expansion:** 10.8 × 10⁻⁶ cm/cm°C (400°C)**Thermal conductivity:** 0.172 w/cm°C (25°C)**Electrical resistivity:** 57 × 10⁻⁶ ohm-cm (25°C)**Ionization potential (1st):** 6.38 eV**Electron work function ϕ :** 3.1 eV**Oxidation potential:** Y → Y³⁺ + 3e = 2.372 V**Chemical valence:** 3**Electrochemical equivalents:** 1.1057 g/amp-hr**Ionic radius:** 0.900 Å (Y³⁺)**Valence electron potential (-eV):** 48.0**Principal quantum number:** 5**Principal electron shells:** K L M N O**Electronic configuration:** 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d¹ 5s²**Valence electrons:** 4d¹ 5s²**Crystal form:** Hexagonal, close packed**Cross section σ :** 1.3 ± 0.1 barns**Vapor pressure:** 5.31 Pa (at melting point)

Zn

Zinc

30

65.38

IIB

65.38 Zn 30
112.41 Cd 48
200.59 Hg 80

Zinco

Zinc

Zink

Zinc

цинк

𨨗

鋅 亜鉛

Naturally occurring isotopes: 64, 66, 68, 67, 70

Density: 7.133 g/cm³ (25°C)

Melting point: 419.58°C **Boiling point:** 907°C

Latent heat of fusion: 113.0 J/g

Specific heat: 0.388 J/g/°C (25°C)

Coefficient of lineal thermal expansion: 39.7 × 10⁻⁶ cm/cm/°C (20°C)

Thermal conductivity: 1.16 w/cm/°C (25°C)

Electrical resistivity: 5.916 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 9.394 eV

Electron work function ϕ : 4.33 eV

Oxidation potential: Zn → Zn²⁺ + 2e = 0.7628 V

Chemical valence: 2

Electrochemical equivalents: 1.220 g/amp-hr

Ionic radius: 0.740 Å (Zn²⁺)

Valence electron potential (-eV): 38.9

Principal quantum number: 4

Principal electron shells: K L M N

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s²

Valence electrons: 4s²

Crystal form: Hexagonal, close packed

Cross section σ : 1.10 ± 0.04 barns

Vapor pressure: 19.2 Pa (at melting point)

Zr

Zirconium

40

91.22

IVB

47.90
Ti
22
91.22
Zr
40
178.49
Hf
72
104

Zircônio
Zirconium
Zirkonium
Zirconio
цирконий
צירקוניום
鋳 ジルコニウム

Naturally occurring isotopes: 90, 94, 92, 91, 96

Density: 6.506 g/cm³ (20°C)

Melting point: 1852 ± 2°C **Boiling point:** 4377°C

Latent heat of fusion: 251.2 J/g

Specific heat: 0.278 J/g°C (25°C)

Coefficient of lineal thermal expansion: 5.85 × 10⁻⁶ cm/cm°C (20°C)

Thermal conductivity: 0.227 w/cm°C (27°C)

Electrical resistivity: 40 × 10⁻⁶ ohm-cm (20°C)

Ionization potential (1st): 6.84 eV

Electron work function ϕ : 4.05 eV

Oxidation potential: $Zr \rightarrow Zr^{4+} + 4e = 1.529 V$

Chemical valence: 1, 2, 3, 4

Electrochemical equivalents: 0.8509 g/amp-hr

Ionic radius: 0.72 Å (Zr⁴⁺)

Valence electron potential (-eV): 80

Principal quantum number: 5

Principal electron shells: K L M N O

Electronic configuration: 1s² 2s² 2p⁶ 3s² 3p⁶ 3d¹⁰ 4s² 4p⁶ 4d² 5s²

Valence electrons: 4d² 5s²

Crystal form: Hexagonal, close packed

Cross section σ : 0.182 ± 0.005 barns

Vapor pressure: 1.68 × 10⁻³ Pa (at melting point)

Element
104

Kurchatovium Rutherfordium

104

261

IVB

47.90 Ti 22
91.22 Zr 40
178.49 Hf 72
104

Naturally occurring isotopes: None

Chemical valence: 4

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Electronic configuration: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^2 4p^6 4d^{10} 4f^{14} 5s^2 5p^6$
 $5d^{10} 5f^{14} 6s^2 6p^6 6d^2 7s^2$

Valence electrons: $6d^2 7s^2$

Half life: ~65 seconds

Element

105

Nielsbohrium Hahnium

105

(262)

VB

50 9415
V
23
92 9064
Nb
41
180 9479
Ta
73
105

Naturally occurring isotopes: None

Chemical valence: (5)

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Half life: ~40 seconds

Element

106

106

(263)

VIB

51.996 Cr 24
95.94 Mo 42
183.85 W 74
106

Naturally occurring isotopes: None

Chemical valence: (6)

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Half life: ~1 second

Element

107

107

(262)

VII B

54.9380
Mn 25
96.906
Tc 43
186.2
Re 75
107

Naturally occurring isotopes: None

Chemical valence: (7)

Principal quantum number: 7

Principal electron shells: K L M N O P Q

Bibliography

Of the numerous references employed in preparation of the third edition of the "Handbook of the Elements," the following are among the most prominent:

- A. E. Bailey et al., Eds., "Tables of Physical and Chemical Constants," 14th ed., Longman Group Limited, London (1973).
- I. Barin and O. Knacke, "Thermochemical Properties of Inorganic Substances," Springer-Verlag, Berlin (1973).
- G. Charlot et al., "Selected Constants: Oxidation-Reduction Potentials of Inorganic Substances in Aqueous Solution," Butterworths, London (1971).
- J. A. Dean, Ed., "Lange's Handbook of Chemistry," 12th ed., McGraw-Hill Book Company, New York (1978).
- S. Fraga and J. Karwowski, "Handbook of Atomic Data," Elsevier Scientific Publishing Company, Amsterdam (1976).
- "Gmelin Handbuch der Anorganischen Chemie," Springer-Verlag, Berlin (up to and including 1980).
- M. Grayson, Executive Ed., "Kirk-Othmer Encyclopedia of Chemical Technology," 3rd ed., Volumes 1-20 incl., John Wiley & Sons, New York (1978-1982 incl.)
- G. B. Naumov et al., Eds., "Handbook of Thermodynamic Data," U.S.S.R. Academy of Sciences, Leningrad (1971).
- L. Pauling, "The Nature of the Chemical Bond," 3rd ed., Cornell University Press, Ithaca, New York (1960).
- G. V. Samsonov, "Handbook of the Physicochemical Properties of the Elements," Plenum, New York (1968).
- R. C. Weast, Editor-in-Chief, "CRC Handbook of Chemistry and Physics, 1983-1984," 64th ed., CRC Press, Inc., Boca Raton, Fla. (1983).
- E. Ya. Zandberg and N. L. Ionov, "Surface Ionization," Science Publishing House, Moscow (1969).

In addition to the reference texts, several primary journals and U.S. government publications were employed. The most commonly utilized were:

- Acta Chemica Scandinavica
- Acta Crystallographica
- Analytical Chemistry
- Bulletin of the American Physical Society
- Canadian Journal of Chemistry
- Canadian Journal of Physics
- Chemical and Engineering News
- Chemical Physics Letters
- Chemical Reviews
- Chemische Berichte
- Electrochimica Acta

Helvetica Chimica Acta
Inorganic Chemistry
Journal of American Chemical Society
Journal of Applied Physics
Journal of Chemical and Engineering Data
Journal of Chemical Education
Journal of Chemical Physics
Journal of Inorganic and Nuclear Chemistry
Journal of Less-Common Metals
Journal of Physical and Chemical Reference Data
Journal of Physical Chemistry
Journal of Physics and Chemistry of Solids
Journal of Solid State Chemistry
Journal of the Chemical Society
Journal of the Electrochemical Society
Materials Research Bulletin
Nature
Physical Review
Proceedings of the Physical Society
Proceedings of the Royal Society
Progress in Inorganic Chemistry
Pure and Applied Chemistry
Science
Talanta
Transactions of the Faraday Society